

## ARG44118 anti-NUP155 antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal recognizes NUP155
Tested Reactivity	Hu
Tested Application	FACS, ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	NUP155
Species	Human
Immunogen	Human NUP155 recombinant protein (Position: D133-K715).
Conjugation	Un-conjugated
Alternate Names	NUP155; Nucleoporin 155; KIAA0791; N155; Nuclear Pore Complex Protein Nup155; 155 KDa Nucleoporin; Nucleoporin 155kDa; Nucleoporin Nup155; Nucleoporin 155kD; ATFB15

### Application Instructions

Application table	Application	Dilution
	FACS	1 - 3 µg/10 <sup>6</sup> cells
	ICC/IF	5 µg/ml
	WB	0.1-0.25 µg/ml
Application Note	The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.05% Sodium azide and 4% Trehalose.
Preservative	0.05% Sodium azide
Stabilizer	4% Trehalose
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation


Gene Symbol	NUP155
Gene Full Name	Nucleoporin 155
Background	Nucleoporins are proteins that play an important role in the assembly and functioning of the nuclear pore complex (NPC) which regulates the movement of macromolecules across the nuclear envelope (NE). The protein encoded by this gene plays a role in the fusion of NE vesicles and formation of the double membrane NE. The protein may also be involved in cardiac physiology and may be associated with the pathogenesis of atrial fibrillation. Alternative splicing results in multiple transcript variants of this gene. A pseudogene associated with this gene is located on chromosome 6.
Function	Essential component of nuclear pore complex. Could be essential for embryogenesis. Nucleoporins may be involved both in binding and translocating proteins during nucleocytoplasmic transport.
Calculated Mw	155 kDa
PTM	Disulfide bond, Glycoprotein, Isopeptide bond, Phosphoprotein, Ubl conjugation
Cellular Localization	Membrane, Nuclear pore complex, Nucleus

Images




ARG44118 anti-NUP155 antibody ICC/IF image

Immunofluorescence: A549 stained with ARG44118 anti-NUP155 antibody at 5 µg/ml dilution.



ARG44118 anti-NUP155 antibody WB image

Western blot: 293T, HeLa stained with ARG44118 anti-NUP155 antibody at 0.25 µg/ml dilution.



#### ARG44118 anti-NUP155 antibody FACS image

Flow Cytometry: HeLa stained with ARG44118 anti-NUP155 antibody at  $1\text{ }\mu\text{g}/10^6$  cells dilution.