

# ARG44491 anti-SMIM8 antibody

Package: 50 μg Store at: -20°C

# Summary

Product Description	Rabbit Polyclonal antibody recognizes SMIM8
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Polyclonal
Isotype	lgG
Target Name	SMIM8
Species	Human
Immunogen	Human SMIM8 recombinant protein
Conjugation	Un-conjugated
Alternate Names	SMIM8; Small Integral Membrane Protein 8; DJ102H19.2; C6orf162; DKFZP586E1923; Chromosome 6 Open Reading Frame 162; UPF0708 Protein C6orf162

# **Application Instructions**

Application table	Application	Dilution
	IHC-P	2-5 μg/ml
Application Note	The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


# Properties

Form	Liquid	
Purification	Affinity purified with Immunogen.	
Buffer	0.9% NaCl, 0.2% Na2HPO4, 0.05% Sodium azide and 4% Trehalose.	
Preservative	0.05% Sodium azide	
Stabilizer	4% Trehalose	
Concentration	0.5 mg/ml	
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.	
Note	For laboratory research only, not for drug, diagnostic or other use.	

### Bioinformation


Gene Symbol	SMIM8
Gene Full Name	Small Integral Membrane Protein 8
Background	Predicted to be integral component of membrane.
Calculated Mw	11 kDa
Cellular Localization	Membrane

### Images



# ARG44491 anti-SMIM8 antibody IHC-P image

Immunohistochemistry: Human liver cancer stained with ARG44491 anti-SMIM8 antibody at 2  $\mu g/mL$  dilution.



### ARG44491 anti-SMIM8 antibody IHC-P image

Immunohistochemistry: Human spleen stained with ARG44491 anti-SMIM8 antibody at 2  $\mu\text{g}/\text{mL}$  dilution.



### ARG44491 anti-SMIM8 antibody IHC-P image

Immunohistochemistry: Human testicular seminoma stained with ARG44491 anti-SMIM8 antibody at 2  $\mu\text{g}/\text{mL}$  dilution.