

ARG45646 anti-COPS5 / JAB1 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes COPS5 / JAB1
Tested Reactivity	Hu, Ms, Rat
Tested Application	FACS, ICC/IF, IHC-Fr, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	COPS5 / JAB1
Species	Human
Immunogen	Recombinant protein containing to human COPS5 / JAB1 .
Conjugation	Un-conjugated
Alternate Names	COPS5; COP9 signalosome subunit 5; MOV-34; Jun activation domain-binding protein 1; EC 3.4.-.-; JAB1; Signalosome subunit 5; SGN5; COP9 signalosome complex subunit 5; CSN5

Application Instructions

Application table	Application	Dilution
	FACS	1 - 3 µg/10 ⁶ cells
	ICC/IF	2 µg/ml
	IHC-Fr	0.5-1 µg/ml
	IHC-P	0.5-1 µg/ml
	WB	0.1-0.5 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	38 kDa	

Properties


Form	Liquid
Purification	Affinity purified
Buffer	0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml

Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	COPS5
Gene Full Name	COP9 signalosome subunit 5
Background	The protein encoded by this gene is one of the eight subunits of COP9 signalosome, a highly conserved protein complex that functions as an important regulator in multiple signaling pathways. The structure and function of COP9 signalosome is similar to that of the 19S regulatory particle of 26S proteasome. COP9 signalosome has been shown to interact with SCF-type E3 ubiquitin ligases and act as a positive regulator of E3 ubiquitin ligases. This protein is reported to be involved in the degradation of cyclin-dependent kinase inhibitor CDKN1B/p27Kip1. It is also known to be an coactivator that increases the specificity of JUN/AP1 transcription factors. [provided by RefSeq, Jul 2008]
Function	Probable protease subunit of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of the SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IκappaBalpha/NFKBIA, ITPK1 and IRF8, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. In the complex, it probably acts as the catalytic center that mediates the cleavage of Nedd8 from cullins. It however has no metalloprotease activity by itself and requires the other subunits of the CSN complex. Interacts directly with a large number of proteins that are regulated by the CSN complex, confirming a key role in the complex. Promotes the proteasomal degradation of BRSK2. [UniProt]
Calculated Mw	38 kDa
PTM	Acetylation. [UniProt]
Cellular Localization	Cytoplasm, cytosol. Nucleus. Cytoplasm, perinuclear region. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle. Note=Nuclear localization is diminished in the presence of IFIT3. [UniProt]

Images




ARG45646 anti-COPS5 / JAB1 antibody IHC-P image

Immunohistochemistry: Human mammary cancer stained with ARG45646 anti-COPS5 / JAB1 antibody at 2 µg/ml dilution.




ARG45646 anti-COP55 / JAB1 antibody ICC/IF image

Immunofluorescence: A431 stained with ARG45646 anti-COP55 / JAB1 antibody at 5 $\mu\text{g}/\text{ml}$ dilution.




ARG45646 anti-COP55 / JAB1 antibody WB image

Western blot: A549 stained with ARG45646 anti-COP55 / JAB1 antibody at 0.5 $\mu\text{g}/\text{ml}$ dilution.




ARG45646 anti-COP55 / JAB1 antibody FACS image

Flow Cytometry: U2OS stained with ARG45646 anti-COP55 / JAB1 antibody at 1 $\mu\text{g}/10^6$ cells dilution.




ARG45646 anti-COP55 / JAB1 antibody IHC-P image

Immunohistochemistry: Rat testis stained with ARG45646 anti-COP55 / JAB1 antibody at 2 $\mu\text{g}/\text{ml}$ dilution.




ARG45646 anti-COPS5 / JAB1 antibody WB image

Western blot: Rat brain stained with ARG45646 anti-COPS5 / JAB1 antibody at 0.5 µg/ml dilution.



ARG45646 anti-COPS5 / JAB1 antibody IHC-P image

Immunohistochemistry: Mouse testis stained with ARG45646 anti-COPS5 / JAB1 antibody at 2 µg/ml dilution.



ARG45646 anti-COPS5 / JAB1 antibody WB image

Western blot: Mouse brain stained with ARG45646 anti-COPS5 / JAB1 antibody at 0.5 µg/ml dilution.