

ARG46006 anti-ANAPC13 antibody

Package: 50 µg

Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes ANAPC13
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	ANAPC13
Species	Human
Immunogen	A 17 amino acid synthetic peptide within the last 50 amino acids of human ANAPC13.
Conjugation	Un-conjugated
Alternate Names	ANAPC13; anaphase promoting complex subunit 13; APC13; SWM1; APC13; Anaphase-promoting complex subunit 13; Cycosome subunit 13

Application Instructions

Application table	Application	Dilution
	ICC/IF	Assay-dependent
	WB	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	14 kDa	

Properties

Purification	Affinity chromatography purified
Buffer	PBS and 0.02% Sodium azide.
Preservative	0.02% Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -56°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	ANAPC13
Gene Full Name	anaphase promoting complex subunit 13
Background	This gene encodes a component of the anaphase promoting complex, a large ubiquitin-protein ligase that controls cell cycle progression by regulating the degradation of cell cycle regulators such as B-type cyclins. The encoded protein is evolutionarily conserved and is required for the integrity and ubiquitin ligase activity of the anaphase promoting complex. Pseudogenes and splice variants have been found for this gene; however, the biological validity of some of the splice variants has not been determined. [provided by RefSeq, Nov 2008]
Function	Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle (PubMed:15060174, PubMed:18485873). The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (PubMed:15060174, PubMed:18485873). The APC/C complex catalyzes assembly of branched 'Lys-11'-/Lys-48'-linked branched ubiquitin chains on target proteins (PubMed:29033132). [UniProt]
Calculated Mw	15 kDa
PTM	Ubl conjugation. [UniProt]
Cellular Localization	Nucleus. [UniProt]