

ARG52294 anti-GABAA Receptor alpha 6 antibody

Package: 50 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes GABAA Receptor alpha 6
Tested Reactivity	Ms, Rat
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	GABAA Receptor alpha 6
Species	Rat
Immunogen	Fusion protein from the cytoplasmic loop of the alpha 6 subunit
Conjugation	Un-conjugated
Alternate Names	A; Gamma-aminobutyric acid receptor subunit alpha-6; GABA

Application Instructions

Application table	Application	Dilution
	WB	1:1,000
Application Note	<p>Specific for the ~57k α6-subunit of the GABAA receptor in Western blots. Labeling is absent in α6-subunit knockout animals.</p> <p>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.</p>	

Properties

Form	Liquid
Purification	Affinity Purified
Buffer	10 mM HEPES (pH 7.5), 150 mM NaCl, 0.1 mg/ml BSA and 50% Glycerol
Stabilizer	0.1 mg/ml BSA, 50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 14399 Mouse
----------------	-------------------------------------

[GeneID: 29708 Rat](#)

[Swiss-port # P16305 Mouse](#)

[Swiss-port # P30191 Rat](#)

Gene Symbol

GABRA6

Gene Full Name

gamma-aminobutyric acid (GABA) A receptor, alpha 6

Background

Gamma-aminobutyric acid (GABA) is the primary inhibitory neurotransmitter in the central nervous system, causing a hyperpolarization of the membrane through the opening of a Cl⁻ channel associated with the GABAA receptor (GABAA-R) subtype. GABAA-Rs are important therapeutic targets for a range of sedative, anxiolytic, and hypnotic agents and are implicated in several diseases including epilepsy, anxiety, depression, and substance abuse. The GABAA-R is a multimeric subunit complex. To date six α s, four β s and four γ s, plus alternative splicing variants of some of these subunits, have been identified (Olsen and Tobin, 1990; Whiting et al., 1999; Ogris et al., 2004). Injection in oocytes or mammalian cell lines of cRNA coding for α - and β -subunits results in the expression of functional GABAA-Rs sensitive to GABA. However, coexpression of a γ -subunit is required for benzodiazepine modulation. The various effects of the benzodiazepines in brain may also be mediated via different α - subunits of the receptor (McKernan et al., 2000; Mehta and Ticku, 1998; Ogris et al., 2004; Pölzl et al., 2003).


Research Area

Neuroscience antibody

Calculated Mw

51 kDa

Images



ARG52294 anti-GABAA Receptor alpha 6 antibody WB image

Western Blot: 5-7 ug of mouse cerebellum lysates from wild type (control) and alpha 6 knockout (alpha 6 K/O) animals showing specific immunolabeling of the ~57k alpha 6-subunit of the GABAA-R in the wild type but not in the alpha 6 K/O animals when stained with GABAA Receptor alpha 6 antibody (ARG52294).