

# ARG55261 anti-LGR5 / GPR49 antibody

Package: 100 μl Store at: -20°C

# Summary

Product Description	Rabbit Polyclonal antibody recognizes LGR5 / GPR49
Tested Reactivity	Hu, Ms, Rat
Tested Application	FACS, ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	lgG
Target Name	LGR5 / GPR49
Species	Human
Immunogen	KLH-conjugated synthetic peptide corresponding to aa. 689-719 of Human LGR5 (Swiss: O75473).
Conjugation	Un-conjugated
Alternate Names	Leucine-rich repeat-containing G-protein coupled receptor 5; G-protein coupled receptor 49; GRP49; G- protein coupled receptor 67; GPR67; GPR49; HG38; G-protein coupled receptor HG38; FEX

### **Application Instructions**

Application table	Application	Dilution
	FACS	1:10 - 1:50
	ICC/IF	1:100
	WB	1:1000
Application Note	* The dilutions indicate recomme should be determined by the scie	nded starting dilutions and the optimal dilutions or concentrations ntist.

## Properties

Form	Liquid
Purification	Purification with Protein A and immunogen peptide.
Buffer	PBS and 0.09% (W/V) Sodium azide
Preservative	0.09% (W/V) Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol Gene Full Name Background	LGR5 leucine-rich repeat containing G protein-coupled receptor 5 The protein encoded by this gene is a leucine-rich repeat-containing receptor (LGR) and member of the G protein-coupled, 7-transmembrane receptor (GPCR) superfamily. The encoded protein is a receptor for R- spondins and is involved in the canonical Wnt signaling pathway. This protein plays a role in the formation and maintenance of adult intestinal stem cells during postembryonic development. Several transcript
Function	Receptor for R-spondins that potentiates the canonical Wnt signaling pathway and acts as a stem cell marker of the intestinal epithelium and the hair follicle. Upon binding to R-spondins (RSPO1, RSPO2, RSPO3 or RSPO4), associates with phosphorylated LRP6 and frizzled receptors that are activated by extracellular Wnt receptors, triggering the canonical Wnt signaling pathway to increase expression of target genes. In contrast to classical G-protein coupled receptors, does not activate heterotrimeric G- proteins to transduce the signal. Involved in the development and/or maintenance of the adult intestinal stem cells during postembryonic development. [UniProt]
Research Area	Cancer antibody; Developmental Biology antibody; Signaling Transduction antibody
Calculated Mw	100 kDa
Cellular Localization	Cell membrane; Multi-pass membrane protein. Golgi apparatus, trans-Golgi network membrane; Multi- pass membrane protein. Note=Rapidly and constitutively internalized to the trans-Golgi network at steady state. Internalization to the trans- Golgi network may be the result of phosphorylation at Ser-861 and Ser-864; however, the phosphorylation event has not been proven (PubMed:23439653).
Images	



#### ARG55261 anti-LGR5 / GPR49 antibody WB image

Western blot: 30  $\mu g$  of HT29, Mouse kidney and Rat kidney lysates stained with ARG55261 anti-LGR5 / GPR49 antibody at 1:1000 dilution.



#### ARG55261 anti-LGR5 / GPR49 antibody FACS image

Flow Cytometry: HepG2 cells stained with ARG55261 anti-LGR5 / GPR49 antibody (bottom histogram) or without primary antibody control (top histogram), followed by incubation with FITC labelled secondary antibody.