

Product datasheet

info@arigobio.com

ARG56232 anti-PACAP Receptor antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes PACAP Receptor

Tested Reactivity Hu, Ms, Rat
Tested Application ICC/IF, WB
Host Rabbit
Clonality Polyclonal

Isotype IgG

Target Name PACAP Receptor

Species Human

Immunogen Synthetic peptide of Human PACAP receptor

Conjugation Un-conjugated

Alternate Names PACAP-R1; PAC1; PACAP type I receptor; PACAP-R-1; Pituitary adenylate cyclase-activating polypeptide

type I receptor; PAC1R; PACAPR; PACAPRI

Application Instructions

Application table	Application	Dilution	
	ICC/IF	1:50 - 1:200	
	WB	1:500 - 1:2000	
Application Note		* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa and Mouse stomach		

Properties

Form Liquid

Purification Affinity purification with immunogen.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol ADCYAP1R1

Gene Full Name adenylate cyclase activating polypeptide 1 (pituitary) receptor type I

Background

This gene encodes type I adenylate cyclase activating polypeptide receptor, which is a membraneassociated protein and shares significant homology with members of the glucagon/secretin receptor
family. This recentor mediates diverse biological actions of adenylate cyclase activating polypeptide 1 and

family. This receptor mediates diverse biological actions of adenylate cyclase activating polypeptide 1 and is positively coupled to adenylate cyclase. Multiple alternatively spliced transcript variants encoding

distinct isoforms have been identified. [provided by RefSeq, Dec 2010]


This is a receptor for PACAP-27 and PACAP-38. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase. May regulate the release of adrenocorticotropin, luteinizing hormone, growth hormone, prolactin, epinephrine, and catecholamine. May play a role in spermatogenesis and

sperm motility. Causes smooth muscle relaxation and secretion in the gastrointestinal tract. [UniProt]

Calculated Mw 53 kDa

Images


Function



ARG56232 anti-PACAP Receptor antibody WB image

Western blot: HeLa cell lysate stained with ARG56232 anti-PACAP Receptor antibody.





Mouse stomach

ARG56232 anti-PACAP Receptor antibody WB image

Western blot: 40 μg of Mouse stomach lysate stained with ARG56232 anti-PACAP Receptor antibody at 1:2000 dilution.