

## ARG56306 anti-PFKM antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes PFKM
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	PFKM
Species	Human
Immunogen	Recombinant protein of Human PFKM
Conjugation	Un-conjugated
Alternate Names	PFK-A; 6-phosphofructokinase type A; PPP1R122; PFKX; ATP-dependent 6-phosphofructokinase, muscle type; EC 2.7.1.11; Phosphofructo-1-kinase isozyme A; PFK1; ATP-PFK; GSD7; PFK-1; Phosphohexokinase; PFK-M; PFKA; Fructose 6 Phosphate Kinase

### Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	22Rv-1	
Observed Size	~ 90 kDa	

### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


## Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	PFKM
Gene Full Name	phosphofructokinase, muscle
Background	Three phosphofructokinase isozymes exist in humans: muscle, liver and platelet. These isozymes function as subunits of the mammalian tetramer phosphofructokinase, which catalyzes the phosphorylation of fructose-6-phosphate to fructose-1,6-bisphosphate. Tetramer composition varies depending on tissue type. This gene encodes the muscle-type isozyme. Mutations in this gene have been associated with glycogen storage disease type VII, also known as Tarui disease. Alternatively spliced transcript variants have been described.[provided by RefSeq, Nov 2009]
Function	Catalyzes the phosphorylation of D-fructose 6-phosphate to fructose 1,6-bisphosphate by ATP, the first committing step of glycolysis. [UniProt]
Calculated Mw	85 kDa
PTM	GlcNAcylation decreases enzyme activity.

## Images



ARG56306 anti-PFKM antibody ICC/IF image

Immunofluorescence: L929 cells stained with ARG56306 anti-PFKM antibody at 1:100 dilution.



22Rv-1

ARG56306 anti-PFKM antibody WB image

Western blot: 22Rv-1 cell lysate stained with ARG56306 anti-PFKM antibody.