

# Product datasheet

info@arigobio.com

# ARG56311 anti-HADHB antibody

Package: 100 μl Store at: -20°C

# **Summary**

Product Description Rabbit Polyclonal antibody recognizes HADHB

Tested Reactivity Hu, Ms, Rat
Tested Application ICC/IF, WB
Host Rabbit
Clonality Polyclonal
Isotype IgG

....

Target Name HADHB
Species Human

Immunogen Recombinant protein of Human HADHB

Conjugation Un-conjugated

Alternate Names MSTP029; ECHB; Trifunctional enzyme subunit beta, mitochondrial; TP-BETA; Beta-ketothiolase; EC

2.3.1.16; Acetyl-CoA acyltransferase; TP-beta; MTPB

# **Application Instructions**

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	SW480	

#### **Properties**

Form Liquid

Purification Affinity purification with immunogen.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol HADHB

Gene Full Name hydroxyacyl-CoA dehydrogenase/3-ketoacyl-CoA thiolase/enoyl-CoA hydratase (trifunctional protein),

beta subunit


Background This gene encodes the beta subunit of the mitochondrial trifunctional protein, which catalyzes the last

three steps of mitochondrial beta-oxidation of long chain fatty acids. The mitochondrial membrane-bound heterocomplex is composed of four alpha and four beta subunits, with the beta subunit catalyzing the 3-ketoacyl-CoA thiolase activity. The encoded protein can also bind RNA and decreases the stability of some mRNAs. The genes of the alpha and beta subunits of the mitochondrial trifunctional protein are located adjacent to each other in the human genome in a head-to-head orientation. Mutations in this gene result in trifunctional protein deficiency. Alternatively spliced transcript variants encoding different

isoforms have been described. [provided by RefSeq, Jul 2013]


Calculated Mw 51 kDa

#### **Images**



#### ARG56311 anti-HADHB antibody ICC/IF image

Immunofluorescence: A549 cells stained with ARG56311 anti-HADHB antibody.



## ARG56311 anti-HADHB antibody WB image

Western blot: SW480 cell lysate stained with ARG56311 anti-HADHB antibody.