

# Product datasheet

info@arigobio.com

ARG56389 anti-LSM2 antibody

Package: 100 μl Store at: -20°C

## Summary

Product Description Rabbit Polyclonal antibody recognizes LSM2

Tested Reactivity Hu

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name LSM2

Species Human

Immunogen Recombinant protein of Human LSM2

Conjugation Un-conjugated

Alternate Names U6 snRNA-associated Sm-like protein LSm2; YBL026W; G7B; snRNP core Sm-like protein Sm-x5; snRNP;

Small nuclear ribonuclear protein D homolog; Protein G7b; C6orf28

### **Application Instructions**

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	NCI-H460	

# **Properties**

Form Liquid

Purification Affinity purification with immunogen.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Database links GeneID: 57819 Human

Swiss-port # Q9Y333 Human

Gene Symbol LSM2

Gene Full Name LSM2 homolog, U6 small nuclear RNA and mRNA degradation associated

Background This gene encodes a member of the LSm family of RNA-binding proteins. LSm proteins form stable

heteromers that bind specifically to the 3'-terminal oligo(U) tract of U6 snRNA and may play a role in pre-mRNA splicing by mediating U4/U6 snRNP formation. Pseudogenes of this gene are located on the


short arm of chromosomes 6 and 19. [provided by RefSeq, Nov 2011]

Function Binds specifically to the 3'-terminal U-tract of U6 snRNA. May be involved in pre-mRNA splicing.

[UniProt]

Calculated Mw 11 kDa

### **Images**



### ARG56389 anti-LSM2 antibody WB image

Western blot: NCI-H460 cell lysate stained with ARG56389 anti-LSM2 antibody.  $\label{eq:local_state} % \begin{subarray}{ll} \end{subarray} % \be$