

## ARG56596 anti-CCL3 / MIP1 alpha antibody [1.2\_3E8-2H6-2B6]

Package: 100 µg, 50 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [1.2_3E8-2H6-2B6] recognizes CCL3 / MIP1 alpha
Tested Reactivity	Hu
Tested Application	ELISA, Neut, WB
Host	Mouse
Clonality	Monoclonal
Clone	1.2_3E8-2H6-2B6
Isotype	IgG1, kappa
Target Name	CCL3 / MIP1 alpha
Species	Human
Immunogen	E.coli derived Recombinant Human MIP-1α (CCL3). (ASLAADTPTA CCFSYTSRQI PQNFIADYFE TSSQCSKPGV IFLTKRSRQV CADPSEEWVQ KYVSDLELSA)
Conjugation	Un-conjugated
Alternate Names	MIP-1-alpha; SIS-beta; PAT 464.1; Tonsillar lymphocyte LD78 alpha protein; G0/G1 switch regulatory protein 19-1; Small-inducible cytokine A3; C-C motif chemokine 3; LD78-alpha; 4-69; Macrophage inflammatory protein 1-alpha; MIP1A; SCYA3; GOS19-1; LD78ALPHA

### Application Instructions

Application table	Application	Dilution
	ELISA	Sandwich: 2.0 - 4.0 µg/ml with ARG56761 as a detection antibody
	Neut	5.0 - 10.0 µg/ml (To yield [ND50] of the biological activity of Human MIP - 1α (100 ng/ml) )
	WB	2.0 - 4.0 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Database links	<a href="#">GeneID: 6348 Human</a> <a href="#">Swiss-port # P10147 Human</a>
Gene Symbol	CCL3
Gene Full Name	chemokine (C-C motif) ligand 3
Background	This locus represents a small inducible cytokine. The encoded protein, also known as macrophage inflammatory protein 1 alpha, plays a role in inflammatory responses through binding to the receptors CCR1, CCR4 and CCR5. Polymorphisms at this locus may be associated with both resistance and susceptibility to infection by human immunodeficiency virus type 1.[provided by RefSeq, Sep 2010]
Function	Monokine with inflammatory and chemokinetic properties. Binds to CCR1, CCR4 and CCR5. One of the major HIV-suppressive factors produced by CD8+ T-cells. Recombinant MIP-1-alpha induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). [UniProt]
Calculated Mw	10 kDa
PTM	N-terminal processed form LD78-alpha(4-69) is produced by proteolytic cleavage after secretion from HTLV1-transformed T-cells.

Images

