

ARG56748 anti-RANTES antibody (Biotin)

Package: 50 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes RANTES
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	RANTES
Species	Human
Immunogen	E.coli derived Recombinant Human RANTES. (SPYSSDTTPC CFAYIARPLP RAHIKEYFYT SGKCSNPVV FVTRKNRQVC ANPEKKWVRE YINSLEMS)
Conjugation	Biotin
Alternate Names	3-68; C-C motif chemokine 5; SIS-delta; D17S136E; SISd; T cell-specific protein P228; EoCP; Small-inducible cytokine A5; Eosinophil chemotactic cytokine; TCP228; 4-68; RANTES; SCYA5; eoCP; T-cell-specific protein RANTES


Application Instructions

Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG56638 as a capture antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties


Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Database links	GeneID: 6352 Human Swiss-port # P13501 Human
Gene Symbol	CCL5
Gene Full Name	chemokine (C-C motif) ligand 5
Background	<p>This gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, functions as a chemoattractant for blood monocytes, memory T helper cells and eosinophils. It causes the release of histamine from basophils and activates eosinophils. This cytokine is one of the major HIV-suppressive factors produced by CD8+ cells. It functions as one of the natural ligands for the chemokine receptor chemokine (C-C motif) receptor 5 (CCR5), and it suppresses in vitro replication of the R5 strains of HIV-1, which use CCR5 as a coreceptor. Alternative splicing results in multiple transcript variants that encode different isoforms. [provided by RefSeq, Jul 2013]</p>
Function	<p>Chemoattractant for blood monocytes, memory T-helper cells and eosinophils. Causes the release of histamine from basophils and activates eosinophils. May activate several chemokine receptors including CCR1, CCR3, CCR4 and CCR5. One of the major HIV-suppressive factors produced by CD8+ T-cells. Recombinant RANTES protein induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). The processed form RANTES(3-68) acts as a natural chemotaxis inhibitor and is a more potent inhibitor of HIV-1-infection. The second processed form RANTES(4-68) exhibits reduced chemotactic and HIV-suppressive activity compared with RANTES(1-68) and RANTES(3-68) and is generated by an unidentified enzyme associated with monocytes and neutrophils. May also be an agonist of the G protein-coupled receptor GPR75, stimulating inositol trisphosphate production and calcium mobilization through its activation. Together with GPR75, may play a role in neuron survival through activation of a downstream signaling pathway involving the PI3, Akt and MAP kinases. By activating GPR75 may also play a role in insulin secretion by islet cells. [UniProt]</p>
Highlight	<p>Related products: RANTES antibodies; RANTES ELISA Kits; Anti-Rabbit IgG secondary antibodies; Related news: circNDUFB2, a circular RNA (circRNA), activates anti-tumor immunity</p>
Calculated Mw	10 kDa
PTM	<p>N-terminal processed form RANTES(3-68) is produced by proteolytic cleavage, probably by DPP4, after secretion from peripheral blood leukocytes and cultured sarcoma cells. The identity of the O-linked saccharides at Ser-27 and Ser-28 are not reported in PubMed:1380064. They are assigned by similarity.</p>



ARG56748 anti-RANTES antibody (Biotin) standard curve image

Direct ELISA: ARG56748 anti-RANTES antibody (Biotin) at 0.25 - 1.0 μ g/ml results of a typical standard run with optical density.



ARG56748 anti-RANTES antibody (Biotin) standard curve image

Sandwich ELISA: ARG56748 anti-RANTES antibody (Biotin) as a detection antibody at 0.25 - 1.0 μ g/ml combined with ARG56638 anti-RANTES antibody as a capture antibody. Results of a typical standard run with optical density.