

ARG57535 anti-VSV-G tag antibody

Package: 100 μg Store at: -20°C

Summary


| Product Description | Mouse Monoclonal antibody recognizes VSV-G tag | |
|---------------------|---|--|
| Tested Reactivity | Other | |
| Tested Application | ICC/IF, WB | |
| Specificity | This antibody can recognize C-terminal, internal, and N-terminal VSV-G fusion proteins. | |
| Host | Mouse | |
| Clonality | Monoclonal | |
| Target Name | VSV-G tag | |
| Species | Others | |
| Immunogen | Synthetic peptide. | |
| Conjugation | Un-conjugated | |
| | | |

Application Instructions

| Application table | Application | Dilution | |
|-------------------|-------------|--|--|
| | ICC/IF | 1:1000 | |
| | WB | 1:5000 | |
| Application Note | | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

| Form | Liquid | |
|---------------------|---|--|
| Purification | Fractional precipitation | |
| Buffer | PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol. | |
| Preservative | 0.02% Sodium azide | |
| Stabilizer | 50% Glycerol | |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. | |
| Note | For laboratory research only, not for drug, diagnostic or other use. | |



ARG57535 anti-VSV-G tag antibody ICC/IF image

Immunofluorescence: 293T cells transfected with a VSV-G-tagged protein and stained with ARG57535 anti-VSV-G tag antibody (red) at 1:2000 dilution. DAPI (blue).

-

ARG57535 anti-VSV-G tag antibody WB image

Western blot: 1 μg of VSV-G fusion protein stained with ARG57535 anti-VSV-G tag antibody at 1:5000 dilution.

VSV-G fusion protein