

Product datasheet

info@arigobio.com

ARG57553 anti-ECHS1 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes ECHS1

Tested Reactivity Hu, Ms, Rat
Tested Application ICC/IF, WB
Host Rabbit
Clonality Polyclonal
Isotype IgG

Target Name ECHS1

Species Human

Immunogen Recombinant protein of Human ECHS1.

Conjugation Un-conjugated

Alternate Names Enoyl-CoA hydratase 1; SCEH; Enoyl-CoA hydratase, mitochondrial; EC 4.2.1.17; Short-chain enoyl-CoA

hydratase; ECHS1D

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:1000 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	U87	
Observed Size	31 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol ECHS1

Gene Full Name enoyl CoA hydratase, short chain, 1, mitochondrial

Background The protein encoded by this gene functions in the second step of the mitochondrial fatty acid beta-

oxidation pathway. It catalyzes the hydration of 2-trans-enoyl-coenzyme A (CoA) intermediates to L-3-hydroxyacyl-CoAs. The gene product is a member of the hydratase/isomerase superfamily. It localizes to the mitochondrial matrix. Transcript variants utilizing alternative transcription initiation


sites have been described in the literature. [provided by RefSeq, Jul 2008]

Function Straight-chain enoyl-CoA thioesters from C4 up to at least C16 are processed, although with decreasing

catalytic rate. [UniProt]

Calculated Mw 31 kDa

Images



ARG57553 anti-ECHS1 antibody ICC/IF image

Immunofluorescence: HeLa cells stained with ARG57553 anti-ECHS1 antibody at 1:100 dilution.



ARG57553 anti-ECHS1 antibody WB image

Western blot: 25 μg of U87 cell lysate stained with ARG57553 anti-ECHS1 antibody at 1:1000 dilution.