

ARG57747 anti-BDNF antibody

Package: 100 μl Store at: -20°C

Summary

| Product Description | Rabbit Polyclonal antibody recognizes BDNF |
|---------------------|---|
| Tested Reactivity | Hu, Ms |
| Predict Reactivity | Rat |
| Tested Application | ICC/IF, IHC-P, WB |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | lgG |
| Target Name | BDNF |
| Species | Human |
| Immunogen | Synthetic peptide from Human BDNF. |
| Conjugation | Un-conjugated |
| Alternate Names | Brain-derived neurotrophic factor; Abrineurin; BULN2; ANON2; BDNF |

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|-----------------|
| | ICC/IF | 1:50 - 1:200 |
| | IHC-P | 1:50 - 1:200 |
| | WB | 1:1000 - 1:2000 |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |
| Positive Control | Mouse heart | |
| Observed Size | ~ 28 kDa | |


Properties

| Form | Liquid |
|---------------------|---|
| Purification | Affinity purified. |
| Buffer | PBS (pH 7.4), 150mM NaCl, 0.02% Sodium azide and 50% Glycerol. |
| Preservative | 0.02% Sodium azide |
| Stabilizer | 50% Glycerol |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| | , For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw |

Bioinformation


| Gene Symbol | BDNF |
|-----------------------|---|
| Gene Full Name | brain-derived neurotrophic factor |
| Background | The protein encoded by this gene is a member of the nerve growth factor family. It is induced by cortical neurons, and is necessary for survival of striatal neurons in the brain. Expression of this gene is reduced in both Alzheimer's and Huntington disease patients. This gene may play a role in the regulation of stress response and in the biology of mood disorders. Multiple transcript variants encoding distinct isoforms have been described for this gene. [provided by RefSeq, Jan 2009] |
| Function | During development, promotes the survival and differentiation of selected neuronal populations of the peripheral and central nervous systems. Participates in axonal growth, pathfinding and in the modulation of dendritic growth and morphology. Major regulator of synaptic transmission and plasticity at adult synapses in many regions of the CNS. The versatility of BDNF is emphasized by its contribution to a range of adaptive neuronal responses including long-term potentiation (LTP), long-term depression (LTD), certain forms of short-term synaptic plasticity, as well as homeostatic regulation of intrinsic neuronal excitability. [UniProt] |
| Highlight | Related products: <u>BDNF antibodies;</u> <u>BDNF ELISA Kits;</u> <u>BDNF recombinant proteins;</u> <u>Anti-Rabbit IgG secondary antibodies;</u> Related news: <u>The role of HDGF in tumor angiogenesis</u> |
| Calculated Mw | 28 kDa |
| PTM | The propeptide is N-glycosylated and glycosulfated. |
| | Converted into mature BDNF by plasmin (PLG). [UniProt] |
| Cellular Localization | Extracellular region or secreted. [UniProt] |


Images



ARG57747 anti-BDNF antibody ICC/IF image

Immunofluorescence: HeLa cells stained with ARG57747 anti-BDNF antibody.





ARG57747 anti-BDNF antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human brain stained with ARG57747 anti-BDNF antibody.

ARG57747 anti-BDNF antibody WB image

Western blot: Mouse heart lysate stained with ARG57747 anti-BDNF antibody.