

Product datasheet

info@arigobio.com

ARG58662 anti-ETS2 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes ETS2

Tested Reactivity Hu, Ms, Rat
Tested Application ICC/IF, WB
Host Rabbit
Clonality Polyclonal

Isotype IgG
Target Name ETS2

Species Human

Immunogen Recombinant fusion protein corresponding to aa. 160-350 of Human ETS2 (NP_005230.1).

Conjugation Un-conjugated

Alternate Names Protein C-ets-2; ETS2IT1

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	53 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol ETS2

Gene Full Name v-ets avian erythroblastosis virus E26 oncogene homolog 2

Background This gene encodes a transcription factor which regulates genes involved in development and apoptosis.

The encoded protein is also a protooncogene and shown to be involved in regulation of telomerase. A pseudogene of this gene is located on the X chromosome. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Jan 2012]

Function Transcription factor activating transcription. Binds specifically the DNA GGAA/T core motif (Ets-binding

site or EBS) in gene promoters and stimulates transcription. [UniProt]


Calculated Mw 53 kDa

PTM Phosphorylation by CDK10 at Ser-220 and Ser-225 creates a phosphodegron that targets ETS2 for

proteasomal degradation. [UniProt]

Cellular Localization Nucleus. [UniProt]

Images



ARG58662 anti-ETS2 antibody ICC/IF image

Immunofluorescence: A549 cells stained with ARG58662 anti-ETS2 antibody.