

Product datasheet

info@arigobio.com

ARG58871 anti-ING1 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes ING1

Tested Reactivity Hu

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG
Target Name ING1

Species Human

Immunogen Synthetic peptide of Human ING1.

Conjugation Un-conjugated

Alternate Names p47ING1a; p24ING1c; Inhibitor of growth protein 1; p33ING1b; p33ING1; p47; p33

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Rat brain	
Observed Size	47 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol ING1

Gene Full Name inhibitor of growth family, member 1

Background This gene encodes a tumor suppressor protein that can induce cell growth arrest and apoptosis. The

encoded protein is a nuclear protein that physically interacts with the tumor suppressor protein TP53 and is a component of the p53 signaling pathway. Reduced expression and rearrangement of this gene have been detected in various cancers. Multiple alternatively spliced transcript variants encoding

distinct isoforms have been reported. [provided by RefSeq, Jul 2008]


Function Cooperates with p53/TP53 in the negative regulatory pathway of cell growth by modulating

p53-dependent transcriptional activation. Implicated as a tumor suppressor gene. [UniProt]

Calculated Mw 47 kDa

Cellular Localization Nucleus. [UniProt]

Images



ARG58871 anti-ING1 antibody WB image

Western blot: 25 μg of Rat brain lysate stained with ARG58871 anti-ING1 antibody at 1:500 dilution.