

ARG58898 anti-Olig2 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Olig2
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Olig2
Species	Human
Immunogen	Synthetic peptide derived from Human Olig2.
Conjugation	Un-conjugated
Alternate Names	bHLHb1; Class E basic helix-loop-helix protein 19; Class B basic helix-loop-helix protein 1; Oligo2; bHLHe19; Oligodendrocyte transcription factor 2; BHLHB1; Protein kinase C-binding protein 2; OLIGO2; Protein kinase C-binding protein RACK17; PRKCBP2; RACK17

Application Instructions

Application table	Application	Dilution
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	32kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation


Gene Symbol	OLIG2
Gene Full Name	oligodendrocyte lineage transcription factor 2
Background	This gene encodes a basic helix-loop-helix transcription factor which is expressed in oligodendroglial tumors of the brain. The protein is an essential regulator of ventral neuroectodermal progenitor cell fate. The gene is involved in a chromosomal translocation t(14;21)(q11.2;q22) associated with T-cell acute lymphoblastic leukemia. Its chromosomal location is within a region of chromosome 21 which has been suggested to play a role in learning deficits associated with Down syndrome. [provided by RefSeq, Jul 2008]
Function	Required for oligodendrocyte and motor neuron specification in the spinal cord, as well as for the development of somatic motor neurons in the hindbrain. Cooperates with OLIG1 to establish the pMN domain of the embryonic neural tube. Antagonist of V2 interneuron and of NKX2-2-induced V3 interneuron development (By similarity). [UniProt]
Calculated Mw	32 kDa
Cellular Localization	Nucleus. Cytoplasm. Note=The NLS contained in the bHLH domain could be masked in the native form and translocation to the nucleus could be mediated by interaction either with class E bHLH partner protein or with NKX2-2. [UniProt]

Images



ARG58898 anti-Olig2 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human brain stained with ARG58898 anti-Olig2 antibody.



ARG58898 anti-Olig2 antibody WB image

Western blot: Human oligodendroglioma lysate stained with ARG58898 anti-Olig2 antibody.