

ARG59201
anti-PREB antibodyPackage: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes PREB
Tested Reactivity	Hu
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	PREB
Species	Human
Immunogen	Recombinant protein corresponding to K42-N236 of Human PREB.
Conjugation	Un-conjugated
Alternate Names	Prolactin regulatory element-binding protein; SEC12; Mammalian guanine nucleotide exchange factor mSec12

Application Instructions

Application table	Application	Dilution
	WB	0.1 - 0.5 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	PREB
Gene Full Name	prolactin regulatory element binding
Background	This gene encodes a protein that specifically binds to a Pit1-binding element of the prolactin (PRL) promoter. This protein may act as a transcriptional regulator and is thought to be involved in some of the developmental abnormalities observed in patients with partial trisomy 2p. This gene overlaps the abhydrolase domain containing 1 (ABHD1) gene on the opposite strand. [provided by RefSeq, Jul 2008]
Function	Was first identified based on its probable role in the regulation of pituitary gene transcription. Binds to the prolactin gene (PRL) promoter and seems to activate transcription (By similarity). Guanine nucleotide exchange factor that activates SARA2. Required for the formation of COPII transport vesicles from the ER. [UniProt]
Calculated Mw	45 kDa
Cellular Localization	Endoplasmic reticulum membrane; Single-pass membrane protein. Nucleus. Note=Concentrates at endoplasmic reticulum exit sites. [UniProt]

Images



ARG59201 anti-PREB antibody WB image

Western blot: 50 µg of samples under reducing conditions. HeLa, Human placenta and A549 whole cell lysates stained with ARG59201 anti-PREB antibody at 0.5 µg/ml, overnight at 4°C.