

ARG62713 anti-CD11b antibody [MEM-170] (Functional Grade)

Package: 100 µg
Store at: -20°C

Summary

Product Description	Functional Grade Mouse Monoclonal antibody [MEM-170] recognizes CD11b
Tested Reactivity	Hu
Tested Application	FACS, FuncSt
Specificity	The clone MEM-170 recognizes CD11b antigen (Mac-1α), a 165-170 kDa transmembrane protein type 1 mainly expressed on monocytes, granulocytes and NK-cells. blocks (not completely) Mac-1-mediated neutrophil binding to fibrinogen. HLDA VI., WS Code A054
Host	Mouse
Clonality	Monoclonal
Clone	MEM-170
Isotype	IgG1
Target Name	CD11b
Species	Human
Immunogen	Human granulocytes
Conjugation	Un-conjugated
Alternate Names	MAC1A; CR3A; CR-3 alpha chain; Cell surface glycoprotein MAC-1 subunit alpha; Integrin alpha-M; MAC-1; CD11 antigen-like family member B; Leukocyte adhesion receptor MO1; MO1A; SLEB6; Neutrophil adherence receptor; CD antigen CD11b; CD11B

Application Instructions

Application table	Application	Dilution
	FACS	1 µg/ml
	FuncSt	Assay-dependent
Application Note	Functional studies: The clone MEM-170 blocks (not completely) Mac-1-mediated neutrophil binding to fibrinogen. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Purification with Protein A.
Purification Note	0.2 µm filter sterilized.
Purity	> 95% (by SDS-PAGE)
Buffer	PBS (pH 7.4)

Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 3684 Human Swiss-port # P11215 Human
Gene Symbol	ITGAM
Gene Full Name	integrin, alpha M (complement component 3 receptor 3 subunit)
Background	CD11b (integrin alpha M chain): Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form a leukocyte-specific integrin referred to as macrophage receptor 1 ('Mac-1'), or inactivated-C3b (iC3b) receptor 3 ('CR3'). The alpha M beta 2 integrin is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009]
Function	CD11b: Integrin ITGAM/ITGB2 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PubMed:9558116, PubMed:20008295). It is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin ITGAM/ITGB2 is also a receptor for fibrinogen, factor X and ICAM1. It recognizes P1 and P2 peptides of fibrinogen gamma chain. Regulates neutrophil migration (PubMed:28807980). In association with beta subunit ITGB2/CD18, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed:21193407). May regulate phagocytosis-induced apoptosis in extravasated neutrophils. May play a role in mast cell development. Required with TYROBP/DAP12 in microglia to control production of microglial superoxide ions which promote the neuronal apoptosis that occurs during brain development. [UniProt]
Highlight	Related products: CD11b antibodies ; CD11b Duos / Panels ; Anti-Mouse IgG secondary antibodies ; Related news: New antibody panels and duos for Tumor immune microenvironment Anti-SerpinB9 therapy, a new strategy for cancer therapy
Research Area	MDSC Marker antibody; Myeloid-derived suppressor cell antibody
Calculated Mw	127 kDa



ARG62713 anti-CD11b antibody [MEM-170] (Functional Grade) FACS image

Flow Cytometry: Human peripheral blood cells stained with ARG62713 anti-CD11b antibody [MEM-170] (Functional Grade), followed by incubation with APC labelled Goat anti-Mouse secondary antibody.