

ARG62863 anti-CD45RA antibody [MEM-56]

Package: 100 μg Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [MEM-56] recognizes CD45RA
Tested Reactivity	Hu
Tested Application	CyTOF [®] -candidate, FACS, IHC-P, IP, WB
Specificity	The clone MEM-56 reacts with CD45RA, a 205-220 kDa single chain type I glycoprotein, variant of CD45 (CD45RA isoform). CD45RA is expressed on most of B lymphocytes, resting and native T lymphocytes, medullar thymocytes and monocytes. HLDA IV; WS Code NL 907
Host	Mouse
Clonality	Monoclonal
Clone	MEM-56
Isotype	lgG2b
Target Name	CD45RA
Species	Human
Immunogen	Human thymocytes and T lymphocytes.
Conjugation	Un-conjugated
Alternate Names	LY5; GP180; Receptor-type tyrosine-protein phosphatase C; CD45; L-CA; CD antigen CD45; Leukocyte common antigen; CD45R; LCA; T200; EC 3.1.3.48; B220

Application Instructions

Application table	Application	Dilution
	CyTOF [®] -candidate	Assay-dependent
	FACS	1 - 4 µg/ml
	IHC-P	5 μg/ml
	IP	Assay-dependent
	WB	Assay-dependent
Application Note	IHC-P: No pre-treatment of * The dilutions indicate records should be determined by the second state of t	ommended starting dilutions and the optimal dilutions or concentrations

Properties


Form	Liquid
Purification	Purified from ascites by protein-A affinity chromatography.
Purity	> 95% (by SDS-PAGE)

Buffer	PBS (pH 7.4) and 15 mM Sodium azide
Preservative	15 mM Sodium azide
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GenelD: 5788 Human
	Swiss-port # P08575 Human
Gene Symbol	PTPRC
Gene Full Name	protein tyrosine phosphatase, receptor type, C
Background	CD45 is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq, Jun 2012]
Function	CD45: Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity. (Microbial infection) Acts as a receptor for human cytomegalovirus protein UL11 and mediates binding of UL11 to T-cells, leading to reduced induction of tyrosine phosphorylation of multiple signaling proteins upon T-cell receptor stimulation and impaired T-cell proliferation. [UniProt]
Research Area	Developmental Biology antibody; Immune System antibody; Neuroscience antibody; Signaling Transduction antibody; Mouse Inflammatory Cell Marker antibody; B Cell Marker antibody
Calculated Mw	147 kDa
PTM	Heavily N- and O-glycosylated.

kDa 1 2 3




ARG62863 anti-CD45RA antibody [MEM-56] WB image

Western blot: HUT-78 human cutaneous T cell lymphoma cell stained with 1) anti-human CD45 control antibody, 2) ARG62863 anti-CD45RA antibody [MEM-56] under non-reducing conditions, 3) ARG62863 anti-CD45RA antibody [MEM-56] under reducing conditions.



ARG62863 anti-CD45RA antibody [MEM-56] IHC-P image

Immunohistochemistry: Paraffin-embedded Human tonsil tissue stained with ARG62863 anti-CD45RA antibody [MEM-56] at 5 $\mu g/ml$ dilution.



ARG62863 anti-CD45RA antibody [MEM-56] CyTOF image

CyTOF: Human peripheral blood stained with ARG62863 anti-CD45RA antibody [MEM-56] (163Dy). Singlet cells were gated for data analysis.