

## ARG62882 anti-CD52 antibody [HI186] (FITC)

Package: 100 tests  
Store at: 4°C

### Summary

Product Description	FITC-conjugated Mouse Monoclonal antibody [HI186] recognizes CD52
Tested Reactivity	Hu
Tested Application	FACS
Specificity	The clone HI186 reacts with CD52 (CAMPATH-1), a 21-28 kDa glycoprotein containing a large N-linked carbohydrate moiety; mature CD52 molecule is actually much smaller (approx. 8-9 kDa). CD52 is expressed at high levels on lymphocytes, monocytes/macrophages and in male reproductive tract. HLDA VI; WS Code BP 523 HLDA VI; WS Code T 6T-057
Host	Mouse
Clonality	Monoclonal
Clone	HI186
Isotype	IgG2b
Target Name	CD52
Species	Human
Immunogen	Human tonsil
Conjugation	FITC
Alternate Names	He5; CAMPATH-1 antigen; Human epididymis-specific protein 5; Cambridge pathology 1 antigen; CDW52; Epididymal secretory protein E5; CDW52; CD antigen CD52

### Application Instructions

Application table	Application	Dilution
	FACS	20 µl / 10 <sup>6</sup> cells
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification Note	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
Buffer	PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA
Preservative	15 mM Sodium azide
Stabilizer	0.2% (w/v) high-grade protease free BSA
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

#### Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

### Database links

[GeneID: 1043 Human](#)

[Swiss-port # P31358 Human](#)

### Gene Symbol

CD52

### Gene Full Name

CD52 molecule

### Background

CD52 (CAMPATH-1, HE5) is a highly glycosylated GPI-anchored 21-28 kDa glycopeptide which is present at high levels on lymphocytes, macrophages, epithelial cells of male reproductive tract and mature sperm. Its 12-amino acid backbone carries a complex N-linked carbohydrate moiety, which differs between sperm and leukocyte CD52, as well as the GPI anchor does. CD52 can be acquired by sperm cells from seminal plasma, where it is released by epithelial cells. Although CD52 is not an essential T-cell costimulator, its triggering results in activation of normal human T cells. CD52 is a very good target for antibody/complement-mediated cell lysis.

### Function

May play a role in carrying and orienting carbohydrate, as well as having a more specific role. [UniProt]


### Research Area

Immune System antibody

### Calculated Mw

7 kDa

## Images



ARG62882 anti-CD52 antibody [HI186] (FITC) FACS image

Flow Cytometry: Human peripheral blood stained with ARG62882 anti-CD52 antibody [HI186] (FITC).