

ARG63962 anti-GCDFP15 antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes GCDFP15
Tested Reactivity	Hu
Tested Application	IHC-P, WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	GCDFP15
Species	Human
Immunogen	C-GANKAQDNTRK
Conjugation	Un-conjugated
Alternate Names	Prolactin-inducible protein; GCDFP-15; Prolactin-induced protein; gp17; SABP; Secretory actin-binding protein; GCDFP15; Gross cystic disease fluid protein 15; GPI4

Application Instructions

Application table	Application	Dilution
	IHC-P	2 - 3 µg/ml
	WB	0.3 - 1 µg/ml
Application Note	IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation


Database links	GeneID: 5304 Human Swiss-port # P12273 Human
Gene Symbol	PIP
Gene Full Name	prolactin-induced protein
Research Area	Cancer antibody; Controls and Markers antibody
Calculated Mw	17 kDa

Images



ARG63962 anti-GCDFP15 antibody WB image

Western blot: A431 cell lysate (35 µg protein in RIPA buffer) stained with ARG63962 anti-GCDFP15 antibody at 0.3 µg/ml dilution.



ARG63962 anti-GCDFP15 antibody IHC-P image

Immunohistochemistry: Paraffin embedded Human Breast. (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG63962 anti-GCDFP15 antibody at 2.5 µg/ml dilution followed by AP-staining.