

ARG64337 anti-CADM4 antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes CADM4
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat
Tested Application	IHC-P, WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	CADM4
Species	Human
Immunogen	C-KDERFQLEEFs
Conjugation	Un-conjugated
Alternate Names	IGSF4C; NECL4; Cell adhesion molecule 4; IgSF4C; Necl-4; Nectin-like protein 4; TSLC1-like protein 2; Immunoglobulin superfamily member 4C; TSL2; NECL-4; synCAM4

Application Instructions

Application table	Application	Dilution
	IHC-P	2.5 µg/ml
	WB	0.1 - 0.3 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 199731 Human Swiss-port # Q8NFZ8 Human
Gene Symbol	CADM4
Gene Full Name	cell adhesion molecule 4
Function	Involved in the cell-cell adhesion. Has calcium- and magnesium-independent cell-cell adhesion activity. May have tumor-suppressor activity. [UniProt]
Research Area	Cancer antibody
Calculated Mw	43 kDa
PTM	N-glycosylated.

Images




ARG64337 anti-CADM4 antibody WB image

Western Blot: Human Brain (Cerebellum) lysate (35µg protein in RIPA buffer). A) ARG64337 anti-CADM4 antibody (0.1µg/ml) and B) ARG64404 anti-CADM4 antibody (0.05µg/ml).



ARG64337 anti-CADM4 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human skeletal muscle tissue. Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). The tissue section was stained with ARG64337 anti-CADM4 antibody at 2.5 µg/ml dilution followed by AP-staining.



ARG64337 anti-CADM4 antibody IHC-P image

Immunohistochemistry: paraffin embedded Human Thymus.
(Steamed antigen retrieval with citrate buffer pH 6) stained with
ARG64337 anti-CADM4 antibody at 2.5 µg/ml dilution followed by
AP-staining.