

### ARG64434 anti-APOBEC2 antibody

Package: 100 μg Store at: -20°C

## Summary

Product Description	Goat Polyclonal antibody recognizes APOBEC2
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat
Tested Application	WB
Host	Goat
Clonality	Polyclonal
lsotype	lgG
Target Name	APOBEC2
Species	Human
Immunogen	C-EAASQNGEDLENLDD
Conjugation	Un-conjugated
Alternate Names	ARCD1; Probable C->U-editing enzyme APOBEC-2; ARP1; EC 3.5.4

## **Application Instructions**

Application table	Application	Dilution
	WB	1 - 3 μg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

#### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

# Bioinformation

Database links	GenelD: 10930 Human
	Swiss-port # Q9Y235 Human
Gene Symbol	APOBEC2
Gene Full Name	apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 2
Function	Probable C to U editing enzyme whose physiological substrate is not yet known. Does not display detectable apoB mRNA editing. Has a low intrinsic cytidine deaminase activity. May play a role in the epigenetic regulation of gene expression through the process of active DNA demethylation. [UniProt]
Research Area	Gene Regulation antibody
Calculated Mw	26 kDa

### Images

A B 250kDa 150kDa	ARG64434 anti-APOBEC2 antibody WB image
100kDa	Western Blot: Human Heart lysate (35 $\mu$ g protein in RIPA buffer)
75kDa	with (B) and without (A) blocking with the immunising peptide.
50kDa	stained with ARG64434 anti-APOBEC2 (aa12 - 26) antibody at 1 μg/ml dilution.
37kDa	
25kDa	
20kDa	
15kDa	
Tonea	