

ARG64593 anti-UNC5CL antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes UNC5CL
Tested Reactivity	Hu
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	UNC5CL
Species	Human
Immunogen	KHCAEQPSHARTY
Conjugation	Un-conjugated
Alternate Names	Protein unc-5 homolog C-like; ZU5 and death domain-containing protein; UNC5C-like protein; ZUD; MUXA

Application Instructions

Application table	<table> <tr> <th>Application</th><th>Dilution</th></tr> <tr> <td>WB</td><td>0.3 - 1 µg/ml</td></tr> </table>	Application	Dilution	WB	0.3 - 1 µg/ml
Application	Dilution				
WB	0.3 - 1 µg/ml				
Application Note	<p>WB: Recommend incubate at RT for 1h.</p> <p>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.</p>				


Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 222643 Human Swiss-port # Q8IV45 Human
Gene Symbol	UNC5CL
Gene Full Name	unc-5 family C-terminal like
Function	Inhibits NF-kappa-B-dependent transcription by impairing NF-kappa-B binding to its targets. [UniProt]
Research Area	Cell Biology and Cellular Response antibody; Cell Death antibody
Calculated Mw	58 kDa

Images



ARG64593 anti-UNC5CL antibody WB image

Western Blot: Human Kidney lysate (35 µg protein in RIPA buffer) stained with ARG64593 anti-UNC5CL antibody at 0.3 µg/ml dilution.