

Product datasheet

info@arigobio.com

ARG64638 anti-APOA1BP antibody

Package: 100 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes APOA1BP

Tested Reactivity Hu

Predict Reactivity Ms, Rat, Cow, Dog, Pig

Tested Application WB
Host Goat

Clonality Polyclonal

Isotype IgG

Target Name APOA1BP
Species Human

 Immunogen
 PPYPDTECVYRLQ

 Conjugation
 Un-conjugated

Alternate Names AIBP; YJEFN1; YjeF_N1; AI-BP; Apolipoprotein A-I-binding protein; NAD; P; EC 5.1.99.6; YjeF N-terminal

domain-containing protein 1

Application Instructions

Application table	Application	Dilution
	WB	0.5 - 1.5 μg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Purified from goat serum by antigen affinity chromatography.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links <u>GeneID: 128240 Human</u>

Swiss-port # Q8NCW5 Human

Background The product of this gene interacts with apolipoprotein A-I (apoA-I), the major apolipoprotein of high-

density lipoproteins (HDLs). It is secreted into some bodily fluids, and its synthesis and secretion are stimulated in vitro by incubating cells with apoA-I. The human genome contains related pseudogenes.

[provided by RefSeq, Jul 2008]

Research Area Cell Biology and Cellular Response antibody

Calculated Mw 32 kDa

PTM Undergoes physiological phosphorylation during sperm capacitation, downstream to PKA activation.

Images

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

ARG64638 anti-APOA1BP antibody WB image

Western Blot: Human Heart lysate (35 μg protein in RIPA buffer) stained with ARG64638 anti-APOA1BP antibody at 0.5 $\mu g/ml$ dilution.