

## ARG64867 anti-CLEC16A antibody

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes CLEC16A
Tested Reactivity	Hu
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	CLEC16A
Immunogen	EEAACAEPVGTAE
Conjugation	Un-conjugated
Alternate Names	Gop-1; KIAA0350; Protein CLEC16A; C-type lectin domain family 16 member A

### Application Instructions

Application table	Application	Dilution
	WB	1 - 3 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties


Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Database links	<a href="#">GeneID: 23274 Human</a> <a href="#">Swiss-port # Q2KHT3 Human</a>
Background	This gene encodes a member of the C-type lectin domain containing family. Single nucleotide polymorphisms in introns of this gene have been associated with diabetes mellitus, multiple sclerosis and rheumatoid arthritis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2011]
Research Area	Immune System antibody
Calculated Mw	118 kDa

## Images

---



ARG64867 anti-CLEC16A antibody WB image

Western Blot: Human Liver lysate (35 µg protein in RIPA buffer) stained with ARG64867 anti-CLEC16A (aa1040-53) antibody at 2 µg/ml dilution.