

ARG64993 anti-TAF7L antibody

Package: 100 µg
Store at: -20°C

Summary

| | |
|---------------------|---|
| Product Description | Goat Polyclonal antibody recognizes TAF7L |
| Tested Reactivity | Hu |
| Tested Application | WB |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | TAF7L |
| Species | Human |
| Immunogen | C-NHFQSVLEQLQLQEK |
| Conjugation | Un-conjugated |
| Alternate Names | Transcription initiation factor TFIID 50 kDa subunit; TATA box-binding protein-associated factor 50 kDa; RNA polymerase II TBP-associated factor subunit Q; TAF2Q; Transcription initiation factor TFIID subunit 7-like; Cancer/testis antigen 40; CT40 |

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|---------------|
| | WB | 0.3 - 1 µg/ml |
| Application Note | WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |


Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification | Purified from goat serum by antigen affinity chromatography. |
| Buffer | Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA. |
| Preservative | 0.02% Sodium azide |
| Stabilizer | 0.5% BSA |
| Concentration | 0.5 mg/ml |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |

Bioinformation

| | |
|----------------|--|
| Database links | GeneID: 54457 Human Swiss-port # Q5H9L4 Human |
| Background | This gene is similar to a mouse gene that encodes a TATA box binding protein-associated factor, and shows testis-specific expression. The encoded protein could be a spermatogenesis-specific component of the DNA-binding general transcription factor complex TFIID. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2009] |
| Research Area | Gene Regulation antibody |
| Calculated Mw | 53 kDa |

Images



ARG64993 anti-TAF7L antibody WB image

Western Blot: Human Testis lysate (35 µg protein in RIPA buffer) stained with ARG64993 anti-TAF7L antibody at 1 µg/ml dilution.