

Product datasheet

info@arigobio.com

ARG65077 anti-FOXI3 / Forkhead box I3 antibody

Package: 100 μg, 50 μg

Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes FOXI3 / Forkhead box I3

Tested Reactivity Hu

Tested Application IHC-P

Host Goat

Clonality Polyclonal

Isotype IgG

Target Name FOXI3 / Forkhead box I3

Species Human

 Immunogen
 C-SPTSISEASADTLQ

 Conjugation
 Un-conjugated

Alternate Names Forkhead box protein I3

Application Instructions

Application table	Application	Dilution
	IHC-P	5 - 10 μg/ml
Application Note	IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations.	

* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations

should be determined by the scientist.

Properties

Form Liquid

Purification Purified from goat serum by antigen affinity chromatography.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GeneID: 344167 Human

Swiss-port # A8MTJ6 Human

Gene Symbol FOXI3


Gene Full Name forkhead box I3

Function Possible transcriptional factor. [UniProt]

Research Area Gene Regulation antibody

Calculated Mw 43 kDa

Images



ARG65077 anti-FOXI3 / Forkhead box I3 antibody IHC-P image

Immunohistochemistry: paraffin embedded Human Bronchus. (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG65077 anti-FOXI3 / Forkhead box I3 antibody at 5 $\mu g/ml$ dilution followed by AP-staining.