

Product datasheet

info@arigobio.com

ARG65165 anti-VLCAD antibody

Package: 100 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes VLCAD

Tested Reactivity Hu
Tested Application WB

Specificity This antibody is expected to recognize both reported isoforms (NP_000009.1; NP_001029031.1).

Host Goat

Clonality Polyclonal

Isotype IgG

Target Name VLCAD

Species Human

Immunogen DKSDSHPSDALTRK-C

Conjugation Un-conjugated

Alternate Names Very long-chain specific acyl-CoA dehydrogenase, mitochondrial; EC 1.3.8.9; LCACD; ACAD6; VLCAD

Application Instructions

Application table	Application	Dilution
	WB	1 - 3 μg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	
	should be determined by the scientist.	

Properties

Form Liquid

Purification Purified from goat serum by antigen affinity chromatography.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed.

freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links <u>GeneID: 37 Human</u>

Swiss-port # P49748 Human

Background The protein encoded by this gene is targeted to the inner mitochondrial membrane where it catalyzes

the first step of the mitochondrial fatty acid beta-oxidation pathway. This acyl-Coenzyme A dehydrogenase is specific to long-chain and very-long-chain fatty acids. A deficiency in this gene product reduces myocardial fatty acid beta-oxidation and is associated with cardiomyopathy. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by

RefSeq, Jul 2008]

Research Area Cancer antibody; Cell Biology and Cellular Response antibody; Metabolism antibody; Signaling

Transduction antibody

Calculated Mw 70 kDa

PTM S-nitrosylation at Cys-237 in liver improves catalytic efficiency.

Images

250kDa
150kDa
100kDa
75kDa

Western Blot: Human Heart lysate (35 μg protein in RIPA buffer)
stained with ARG65165 anti-VLCAD antibody at 2 μg/ml dilution.

50kDa
37kDa

25kDa
20kDa