

ARG65545
anti-CD2 antibody [TS1/8]Package: 100 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [TS1/8] recognizes CD2
Tested Reactivity	Hu
Tested Application	CyTOF®-candidate, FACS, FuncSt
Specificity	The mouse monoclonal antibody TS1/8 recognizes CD2, a 50 kDa glycoprotein present on the human peripheral blood T lymphocytes and NK cells; also expressed by all thymocytes. HLDA V; WS Code S025
Host	Mouse
Clonality	Monoclonal
Clone	TS1/8
Isotype	IgG1
Target Name	CD2
Immunogen	Cytotoxic T lymphocytes
Conjugation	Un-conjugated
Alternate Names	T-cell surface antigen T11/Leu-5; LFA-3 receptor; T-cell surface antigen CD2; SRBC; Erythrocyte receptor; CD antigen CD2; T11; Rosette receptor; LFA-2

Application Instructions

Application table	Application	Dilution
	CyTOF®-candidate	Assay-dependent
	FACS	1 - 4 µg/ml
	FuncSt	Assay-dependent
Application Note	Functional studies: Blocking. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties


Form	Liquid
Purification	Purified from cell culture supernatant by protein-A affinity chromatography.
Purity	> 95% (by SDS-PAGE)
Buffer	PBS (pH 7.4) and 15 mM Sodium azide
Preservative	15 mM Sodium azide
Concentration	1 mg/ml

Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 914 Human Swiss-port # P06729 Human
Gene Symbol	CD2
Gene Full Name	CD2 molecule
Background	CD2 (also known as LFA-2) belongs to T lymphocyte glycoproteins of immunoglobulin superfamily. Its interaction with CD58 stabilizes adhesion between T cells and antigen presenting or target cells. Relatively low affinity of CD2 to CD58 (as measured in solution) is compensated within the two-dimensional cell-cell interface to provide tight adhesion. Moreover, T cell activation induces increased CD2 expression and its lateral mobility, making easier contact between CD2 and CD58. Subsequently, T cell activation causes fixation of CD58-CD2 at sites of cell-cell contact, thereby strengthening intercellular adhesion. CD2 deficiency reduces intestinal inflammation and helps to control infection.
Function	CD2 interacts with lymphocyte function-associated antigen (LFA-3) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function. [UniProt]
Highlight	Related products: CD2 antibodies ; CD2 ELISA Kits ; CD2 Duos / Panels ; Anti-Mouse IgG secondary antibodies ; Related news: CyTOF-candidate Antibodies
Research Area	Developmental Biology antibody; Immune System antibody
Calculated Mw	39 kDa

Images



ARG65545 anti-CD2 antibody [TS1/8] FACS image

Flow Cytometry: Human peripheral blood stained with ARG65545 anti-CD2 antibody [TS1/8] at 4 µg/ml dilution, followed by APC-conjugated Goat anti-Mouse antibody.



ARG65545 anti-CD2 antibody [TS1/8] FACS image

Flow Cytometry: Separation of human CD2 positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed). Human peripheral whole blood stained with ARG65545 anti-CD2 antibody [TS1/8] at 4 $\mu\text{g}/\text{ml}$ dilution, followed by APC-conjugated Goat anti-Mouse antibody.