

## ARG66704 anti-TIE1 antibody

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes TIE1
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	TIE1
Species	Human
Immunogen	Synthetic peptide within aa. 820-900 of Human TIE1.
Conjugation	Un-conjugated
Alternate Names	TIE; Tyrosine-protein kinase receptor Tie-1; JTK14; EC 2.7.10.1

### Application Instructions

Application table	Application	Dilution
	IHC-P	1:100 - 1:300
Application Note	IHC-P: Antigen Retrieval: EDTA buffer (pH 8.0) at high pressure and temperature. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS, 0.02% Sodium azide, 50% Glycerol and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol and 0.5% BSA
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol	TIE1
Gene Full Name	tyrosine kinase with immunoglobulin-like and EGF-like domains 1
Background	This gene encodes a member of the tyrosine protein kinase family. The encoded protein plays a critical role in angiogenesis and blood vessel stability by inhibiting angiopoietin 1 signaling through the endothelial receptor tyrosine kinase Tie2. Ectodomain cleavage of the encoded protein relieves inhibition of Tie2 and is mediated by multiple factors including vascular endothelial growth factor. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Nov 2011]
Function	Transmembrane tyrosine-protein kinase that may modulate TEK/TIE2 activity and contribute to the regulation of angiogenesis. [UniProt]
Calculated Mw	125 kDa
PTM	Phosphorylated on tyrosine residues in response to ANGPT1, most likely by TEK/TIE2. [UniProt]
Cellular Localization	Cell membrane; Single-pass type I membrane protein. [UniProt]


## Images

---



ARG66704 anti-TIE1 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human oophoroma tissue. Antigen Retrieval: EDTA buffer (pH 8.0) at high pressure and temperature. The tissue section was stained with ARG66704 anti-TIE1 antibody at 1:100 dilution, overnight at 4°C.



ARG66704 anti-TIE1 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human kidney tissue. Antigen Retrieval: EDTA buffer (pH 8.0) at high pressure and temperature. The tissue section was stained with ARG66704 anti-TIE1 antibody at 1:200 dilution, overnight at 4°C.