

ARG67138 anti-CD63 antibody

Package: 20 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes CD63
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CD63
Species	Human
Immunogen	Synthetic peptide of Human CD63.
Conjugation	Un-conjugated
Alternate Names	CD63; CD63 Molecule; TSPAN30; ME491; MLA1; Ocular Melanoma-Associated Antigen; CD63 Antigen (Melanoma 1 Antigen); Melanoma-Associated Antigen ME491; Tetraspanin-30; Granulophysin; CD63 Antigen; Tspan-30; OMA81H; Limp1; Lysosomal-Associated Membrane Protein 3; Lysosome Integral Membrane Protein 1; Melanoma-Associated Antigen MLA1; LAMP-3

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:200 - 1:1000
	IHC-P	1:100 - 1:300
	WB	1:500 - 1:2000

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Affinity chromatography purified
Buffer	PBS, 0.02% Sodium azide, 0.5% BSA and 50% Glycerol
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA, 50% Glycerol
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed


before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	CD63
Gene Full Name	CD63 Molecule
Background	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms. [provided by RefSeq, Apr 2012]
Function	Functions as a cell surface receptor for TIMP1 and plays a role in the activation of cellular signaling cascades. Plays a role in the activation of ITGB1 and integrin signaling, leading to the activation of AKT, FAK/PTK2 and MAP kinases. Promotes cell survival, reorganization of the actin cytoskeleton, cell adhesion, spreading and migration, via its role in the activation of AKT and FAK/PTK2. Plays a role in VEGFA signaling via its role in regulating the internalization of KDR/VEGFR2. Plays a role in intracellular vesicular transport processes, and is required for normal trafficking of the PMEL luminal domain that is essential for the development and maturation of melanocytes. Plays a role in the adhesion of leukocytes onto endothelial cells via its role in the regulation of SELP trafficking. May play a role in mast cell degranulation in response to Ms4a2/FcεRI stimulation, but not in mast cell degranulation in response to other stimuli. [Uniprot]
Calculated Mw	26 kDa
PTM	Glycoprotein, Lipoprotein, Palmitate. [Uniprot]
Cellular Localization	Cell membrane, Endosome, Lysosome, Membrane, Secreted. [Uniprot]

Images




ARG67138 anti-CD63 antibody IHC-P image

Immunohistochemistry: Human liver cancer stained with ARG67138 anti-CD63 antibody at 1:200 dilution.



ARG67138 anti-CD63 antibody ICC/IF image

Immunofluorescence: A549 stained with ARG67138 anti-CD63 antibody at 1:200 dilution.



ARG67138 anti-CD63 antibody WB image

Western blot: SKOV3 stained with ARG67138 anti-CD63 antibody.