

ARG70036

Human TGF beta 1 recombinant protein (Active) (His-tagged, C-ter)

Package: 100 µg, 20 µg

Store at: -20°C

Summary

Product Description	E. coli expressed, His-tagged (C-ter) Active Human TGF beta 1 recombinant protein
Tested Application	SDS-PAGE
Target Name	TGF beta 1
Species	Human
A.A. Sequence	Ala279 - Ser390
Expression System	E. coli
Activity	Active
Activity Note	Determined by its ability to inhibit the IL-4 dependent proliferation in HT-2 cells. The ED50 for this effect is 5×10^7 IU/mg. Determined by its ability to induce proliferation in MCF-7 cells The ED50 for this effect is 98% (by SDS-PAGE)
Buffer	20 mM sodium citrate and 0.2 M NaCl (pH 4.5)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely. In some experiments, it recommends to add 10 mM HCl when reconstitute lyophilized protein.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	TGFB1
Gene Full Name	transforming growth factor, beta 1
Background	This gene encodes a member of the transforming growth factor beta (TGFB) family of cytokines, which are multifunctional peptides that regulate proliferation, differentiation, adhesion, migration, and other functions in many cell types. Many cells have TGFB receptors, and the protein positively and negatively regulates many other growth factors. The secreted protein is cleaved into a latency-associated peptide (LAP) and a mature TGFB1 peptide, and is found in either a latent form composed of a TGFB1 homodimer, a LAP homodimer, and a latent TGFB1-binding protein, or in an active form composed of a TGFB1 homodimer. The mature peptide may also form heterodimers with other TGFB family members. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease.[provided by RefSeq, Oct 2009]
Function	Multifunctional protein that controls proliferation, differentiation and other functions in many cell types. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other growth factors. It plays an important role in bone remodeling as it is a potent stimulator of osteoblastic bone formation, causing chemotaxis, proliferation and differentiation in committed osteoblasts. Can promote either T-helper 17 cells (Th17) or regulatory T-cells (Treg) lineage differentiation in a concentration-dependent manner. At high concentrations, leads to FOXP3-mediated suppression of RORC and down-regulation of IL-17 expression, favoring Treg cell development. At low concentrations in concert with IL-6 and IL-21, leads to expression of the IL-17 and IL-23 receptors, favoring differentiation to Th17 cells. [UniProt]

PTM


Glycosylated.

The precursor is cleaved into mature TGF-beta-1 and LAP, which remains non-covalently linked to mature TGF-beta-1 rendering it inactive. [UniProt]

Cellular Localization

Secreted, extracellular space, extracellular matrix. [UniProt]

Images



Human TGF beta 1
recombinant protein

ARG70036 Human TGF beta 1 recombinant protein (Active) (His-tagged, C-ter) SDS-PAGE image

SDS-PAGE analysis of ARG70036 Human TGF beta 1 recombinant protein (Active) (His-tagged, C-ter).