

ARG70057 Human IL13 recombinant protein (Active) (His-tagged, C-ter)

Package: 100 µg, 20 µg
Store at: -20°C

Summary


Product Description	E. coli expressed, His-tagged (C-ter) Active Human IL13 recombinant protein
Tested Application	SDS-PAGE
Target Name	IL13
Species	Human
A.A. Sequence	Gly35 - Asn146
Expression System	E. coli
Activity	Active
Alternate Names	Interleukin-13; P600; IL-13

Properties

Form	Powder
Purification Note	Endotoxin level is < 0.01 EU/µg of the protein, as determined by the LAL test.
Purity	> 95% (by SDS-PAGE)
Buffer	PBS (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	IL13
Gene Full Name	interleukin 13
Background	This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen-induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]
Function	Cytokine. Inhibits inflammatory cytokine production. Synergizes with IL2 in regulating interferon-gamma synthesis. May be critical in regulating inflammatory and immune responses. [UniProt]
Cellular Localization	Secreted. [UniProt]



ARG70057 Human IL13 recombinant protein (Active) (His-tagged, C-ter) SDS-PAGE image

SDS-PAGE analysis of ARG70057 Human IL13 recombinant protein (Active) (His-tagged, C-ter).