

**ARG70064**  
**Human IL20 recombinant protein (Active) (His-tagged, C-ter)**Package: 100 µg, 20 µg  
Store at: -20°C

### Summary


Product Description	E. coli expressed, His-tagged (C-ter) Active Human IL20 recombinant protein
Tested Application	SDS-PAGE
Target Name	IL20
Species	Human
A.A. Sequence	Leu25 - Glu176
Expression System	E. coli
Activity	Active
Activity Note	Determined by its ability to chemoattract BaF3 cells transfected with human IL-20R alpha and IL-20R beta. The ED50 for this effect is < 0.2 ng/mL.
Alternate Names	Interleukin-20; IL-20; IL10D; Cytokine Zcyto10; ZCYTO10

### Properties

Form	Powder
Purification Note	Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Buffer	PBS (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol	IL20
Gene Full Name	interleukin 20
Background	The protein encoded by this gene is a cytokine structurally related to interleukin 10 (IL10). This cytokine has been shown to transduce its signal through signal transducer and activator of transcription 3 (STAT3) in keratinocytes. A specific receptor for this cytokine is found to be expressed in skin and upregulated dramatically in psoriatic skin, suggesting a role for this protein in epidermal function and psoriasis. [provided by RefSeq, Jul 2008]
Function	Proinflammatory and angiogenic cytokine that may be involved in epidermal function and psoriasis. Angiogenic and proliferative activities are antagonized by IL10. May act through STAT3. [UniProt]
Cellular Localization	Secreted. [UniProt]



ARG70064 Human IL20 recombinant protein (Active) (His-tagged, C-ter) SDS-PAGE image

SDS-PAGE analysis of ARG70064 Human IL20 recombinant protein (Active) (His-tagged, C-ter).