

ARG70074

Human IL29 / IFN lambda recombinant protein (Active) (His-tagged, C-ter)

Package: 100 µg, 20 µg

Store at: -20°C

Summary

Product Description	E. coli expressed, His-tagged (C-ter) Active Human IL29 / IFN lambda recombinant protein
Tested Application	SDS-PAGE
Target Name	IL29 / IFN lambda
Species	Human
A.A. Sequence	Pro23 - Thr200
Expression System	E. coli
Activity	Active
Activity Note	Determined by its ability to induce IL-8 secretion in HuH7 cells. The ED50 for this effect is < 6 ng/mL.
Alternate Names	IFN-lambda-1; IL-29; Interferon lambda-1; Interleukin-29; Cytokine Zcyto21; IL29

Properties

Form	Powder
Purification Note	Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Buffer	PBS (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	IFNL1
Gene Full Name	interferon, lambda 1
Background	This gene encodes a cytokine distantly related to type I interferons and the IL-10 family. This gene, interleukin 28A (IL28A), and interleukin 28B (IL28B) are three closely related cytokine genes that form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by the three genes can be induced by viral infection. All three cytokines have been shown to interact with a heterodimeric class II cytokine receptor that consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA). [provided by RefSeq, Jul 2008]
Function	Cytokine with antiviral, antitumour and immunomodulatory activities. Plays a critical role in the antiviral host defense, predominantly in the epithelial tissues. Acts as a ligand for the heterodimeric

class II cytokine receptor composed of IL10RB and IFNLR1, and receptor engagement leads to the activation of the JAK/STAT signaling pathway resulting in the expression of IFN-stimulated genes (ISG), which mediate the antiviral state. Has a restricted receptor distribution and therefore restricted targets: is primarily active in epithelial cells and this cell type-selective action is because of the epithelial cell-specific expression of its receptor IFNLR1. Exerts an immunomodulatory effect by up-regulating MHC class I antigen expression. [UniProt]

Highlight

Related products:

[IL29 antibodies](#); [IL29 ELISA Kits](#); [IL29 recombinant proteins](#);


Related news:

[HMGB1, a biomarker and therapeutic target in COVID-19](#)

Cellular Localization

Secreted. [UniProt]

Images



Human IL29 / IFN lambda
recombinant protein

ARG70074 Human IL29 / IFN lambda recombinant protein (Active)
(His-tagged, C-ter) SDS-PAGE image

SDS-PAGE analysis of ARG70074 Human IL29 / IFN lambda
recombinant protein (Active) (His-tagged, C-ter).