

ARG70114

Human FGF8 (isoform A) recombinant protein (Active) (His-tagged, C-ter)

Package: 100 µg, 20 µg

Store at: -20°C

Summary

Product Description	E. coli expressed, His-tagged (C-ter) Active Human FGF8 (isoform A) recombinant protein
Tested Application	SDS-PAGE
Target Name	FGF8 (isoform A)
Species	Human
A.A. Sequence	Gln52 - Arg223
Expression System	E. coli
Activity	Active
Activity Note	Determined by its ability to induce 3T3 cells proliferation. The ED50 for this effect is < 2 µg/mL.
Alternate Names	Fibroblast growth factor 8; HH6; AIGF; FGF-8; HBGF-8; KAL6; Heparin-binding growth factor 8; Androgen-induced growth factor

Properties

Form	Powder
Purification Note	Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Buffer	PBS (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	FGF8
Gene Full Name	fibroblast growth factor 8 (androgen-induced)
Background	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein is known to be a factor that supports androgen and anchorage independent growth of mammary tumor cells. Overexpression of this gene has been shown to increase tumor growth and angiogenesis. The adult expression of this gene is restricted to testes and ovaries. Temporal and spatial pattern of this gene expression suggests its function as an embryonic epithelial factor. Studies of the mouse and chick homologs revealed roles in midbrain and limb development,

organogenesis, embryo gastrulation and left-right axis determination. The alternative splicing of this gene results in four transcript variants. [provided by RefSeq, Jul 2008]


Function

Plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. Required for normal brain, eye, ear and limb development during embryogenesis. Required for normal development of the gonadotropin-releasing hormone (GnRH) neuronal system. Plays a role in neurite outgrowth in hippocampal cells. [UniProt]

Cellular Localization

Secreted. [UniProt]

Images



ARG70114 Human FGF8 (isoform A) recombinant protein (Active)
(His-tagged, C-ter) SDS-PAGE image

SDS-PAGE analysis of ARG70114 Human FGF8 (isoform A)
recombinant protein (Active) (His-tagged, C-ter).