

ARG70158 Mouse IL6 recombinant protein (Active) (His-tagged, C-ter)

Package: 100 µg, 20 µg
Store at: -20°C

Summary

Product Description	E. coli expressed, His-tagged (C-ter) Active Mouse IL6 recombinant protein
Tested Application	SDS-PAGE
Target Name	IL6
Species	Mouse
A.A. Sequence	Phe25 - Thr211
Expression System	E. coli
Activity	Active
Alternate Names	B-cell stimulatory factor 2; CDF; HSF; BSF-2; Interferon beta-2; IL-6; IFNB2; CTL differentiation factor; Interleukin-6; HGF; Hybridoma growth factor; BSF2; IFN-beta-2

Properties

Form	Powder
Purification Note	Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Buffer	PBS (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	IL6
Gene Full Name	interleukin 6
Background	This gene encodes a cytokine that functions in inflammation and the maturation of B cells. In addition, the encoded protein has been shown to be an endogenous pyrogen capable of inducing fever in people with autoimmune diseases or infections. The protein is primarily produced at sites of acute and chronic inflammation, where it is secreted into the serum and induces a transcriptional inflammatory response through interleukin 6 receptor, alpha. The functioning of this gene is implicated in a wide variety of inflammation-associated disease states, including susceptibility to diabetes mellitus and systemic juvenile rheumatoid arthritis. [provided by RefSeq, Jun 2011]
Function	Cytokine with a wide variety of biological functions. It is a potent inducer of the acute phase response. Plays an essential role in the final differentiation of B-cells into Ig-secreting cells Involved in lymphocyte and monocyte differentiation. Acts on B-cells, T-cells, hepatocytes, hematopoietic progenitor cells and cells of the CNS. Required for the generation of T(H)17 cells. Also acts as a myokine. It is discharged into the bloodstream after muscle contraction and acts to increase the breakdown of fats and to improve

insulin resistance. It induces myeloma and plasmacytoma growth and induces nerve cells differentiation. [UniProt]

Highlight

Related products:

[IL6 antibodies](#); [IL6 ELISA Kits](#); [IL6 recombinant proteins](#);

Related news:

[HMGB1 in inflammation](#)

[Inflammatory Cytokines](#)


PTM

N- and O-glycosylated. [UniProt]

Cellular Localization

Secreted. [UniProt]

Images



Mouse IL6
recombinant protein

ARG70158 Mouse IL6 recombinant protein (Active) (His-tagged, C-ter) SDS-PAGE image

SDS-PAGE analysis of ARG70158 Mouse IL6 recombinant protein (Active) (His-tagged, C-ter).