

Product datasheet

info@arigobio.com

ARG70315
Mouse IL21R recombinant protein (His-tagged, C-ter)

Package: 100 μg Store at: -20°C

Summary

Product Description HEK293 expressed, His-tagged (C-ter) Mouse IL21R recombinant protein.

Tested Reactivity Ms

Tested Application Binding, SDS-PAGE

Target Name IL21R

Species Mouse

A.A. Sequence Cys20 - Pro236 of Mouse IL21R (NP_068687.1) with 6X His tag at the C - terminus.

Expression System HEK293

Alternate Names Interleukin-21 receptor; Novel interleukin receptor; CD360; CD antigen CD360; NILR; IL-21R; IL-21

receptor

Application Instructions

Application Note Binding activity test: Measured by its binding ability in a functional ELISA. Immobilized Recombinant

Mouse IL21R at 2μg/ml (100μl/well) can bind Recombinant Human IL-21 with a linear range of

0.128-83.57 ng/ml.

Properties

Form Powder

Purification Note 0.22 μm filter sterilized. Endotoxin level is 97% (by SDS-PAGE)

Buffer PBS (pH 7.4)

 $\label{eq:Reconstitution} \textbf{Reconstitute to a concentration of 0.1 - 0.5 mg/ml in sterile distilled water.}$

Storage instruction For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and

store at -20°C for up to one month, at 2-8°C for up to one week. Storage in frost free freezers is not

recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol IL21R

Gene Full Name interleukin 21 receptor

Background The protein encoded by this gene is a cytokine receptor for interleukin 21 (IL21). It belongs to the type I

cytokine receptors, and has been shown to form a heterodimeric receptor complex with the common gamma-chain, a receptor subunit also shared by the receptors for interleukin 2, 4, 7, 9, and 15. This receptor transduces the growth promoting signal of IL21, and is important for the proliferation and differentiation of T cells, B cells, and natural killer (NK) cells. The ligand binding of this receptor leads to the activation of multiple downstream signaling molecules, including JAK1, JAK3, STAT1, and STAT3. Knockout studies of a similar gene in mouse suggest a role for this gene in regulating immunoglobulin

www.arigobio.com arigo.nuts about antibodies 1/2

production. Three alternatively spliced transcript variants have been described. [provided by RefSeq, Jul

2010]

Function This is a receptor for interleukin-21. [UniProt]

Calculated Mw 59 kDa


PTM C-mannosylated at Trp-214 in the WSXWS motif, the sugar chain makes extensive hydrogen bonds with

Asn-73 sugar, and bridges the two fibronectin domains transforming the V-shaped receptor into an A-

frame. [UniProt]

Cellular Localization Membrane; Single-pass type I membrane protein. [UniProt]

Images



ARG70315 Mouse IL21R recombinant protein (ECD) (His-tagged, Cter) SDS-PAGE image

SDS-PAGE analysis of ARG70315 Mouse IL21R recombinant protein (ECD) (His-tagged, C-ter).