

Product datasheet

info@arigobio.com

ARG70379
Human IL38 recombinant protein (His-tagged, C-ter)

Package: 100 μ g, 20 μ g

Store at: -20°C

Summary

Product Description E. coli expressed, His-tagged (C-ter) Human IL38 recombinant protein

Tested Application SDS-PAGE

Target Name IL38

Species Human

A.A. Sequence Met1 - Trp152

Expression System E. coli

Alternate Names IL1F10; Interleukin 1 Family Member 10; IL-1HY2; IL1-Theta; IL-1F10; FKSG75; Interleukin-1 Receptor

Antagonist FKSG75; Interleukin-1 Family Member 10; Family Of Interleukin 1-Theta; Interleukin-1 HY2;

Interleukin-38; FIL1 Theta; IL-1 Theta; MGC119832; MGC119833; MGC11983; IL1HY2; IL-38;

Interleukin-1 Receptor Antagonist-Like FIL1 Theta; Interleukin 1 Family, Member 10 (Theta); Interleukin 1 Family Member 10 (Theta); IL-1F10 (Canonical Form IL-1F10a); Interleukin-1 Theta; FIL1- Theta;

FIL1-Theta; FIL1T; IL38

Properties

Form Powder

Purification Note Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.

Purity > 98% (by SDS-PAGE)

Buffer PBS (pH 7.4)

Reconstitution It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less

than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure

the protein is dissolved completely.

Storage instruction For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and

store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol IL1F10

Gene Full Name Interleukin 1 Family Member 10

Background The protein encoded by this gene is a member of the interleukin 1 cytokine family. This gene and eight

other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. This cytokine is thought to participate in a network of interleukin 1 family members to regulate adapted and innate immune responses. Two alternatively spliced transcript variants encoding the same protein have been

reported.


Function Cytokine with immunomodulatory activity. Alone, does not induce cytokine production, but reduces

IL22 and IL17A production by T-cells in response to heat-killed Candida albicans. Reduces IL36G-induced production of IL8 by peripheral blood mononuclear cells. Increases IL6 production by dendritic cells

stimulated by bacterial lipopolysaccharides (LPS). Ligand for IL-36R/IL1RL2.

www.arigobio.com arigo.nuts about antibodies 1/2

Images



ARG70379 Human IL38 recombinant protein (His-tagged, C-ter) SDS-PAGE image

SDS-PAGE analysis of ARG70379 Human IL38 recombinant protein (His-tagged, C-ter) $\,$