

## ARG80823 Human Adiponectin ELISA Kit

Package: 96 wells  
Store at: 4°C

### Summary

Product Description	ARG80823 Human Adiponectin ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human Adiponectin in serum and plasma (heparin).
Tested Reactivity	Hu
Tested Application	ELISA
Target Name	Adiponectin
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm
Sensitivity	< 0.27 ng/ml
Sample Type	Serum and plasma (heparin).
Standard Range	2 - 100 ng/ml
Sample Volume	100 µl
Alternate Names	Adipose most abundant gene transcript 1 protein; ADPN; APM-1; Gelatin-binding protein; APM1; ACDC; Adiponectin; apM-1; ACRP30; ADIPQTL1; GBP28; 30 kDa adipocyte complement-related protein; Adipocyte, C1q and collagen domain-containing protein; Adipocyte complement-related 30 kDa protein

### Application Instructions

Assay Time	1 h, 30 min (RT/shaking), 15 min (RT)
------------	---------------------------------------

### Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Database links	<a href="#">GeneID: 9370 Human</a> <a href="#">Swiss-port # Q15848 Human</a>
Gene Symbol	ADIPOQ
Gene Full Name	adiponectin, C1Q and collagen domain containing
Background	Adiponectin is a 30kDa protein which percentage in serum proteins is 0.01%. It is mainly synthesized by Adipocytes, but also muscle cells and hepatocytes have the ability to synthesize Adiponectin. Until now, IGF-I is the only known natural inductor of the synthesis. It consists of a Collagen-like N-terminal and a globular C-terminal domain. In vivo Adiponectin appears with different oligomers. Beside the trimer

and dimer also high molecular multimers exist. Up to now two different receptors are known, both receptors are ubiquitarily expressed, though the distribution in the tissues varies. The Adiponectin Receptor 1 (AdipoR1) is especially in muscle- and AdipoR2 in liver tissue synthesized.

The significance for the human organism is not clear until now. First studies show, that adiponectin correlates negatively with BMI and thus it could have relevance for the energy metabolism for example through the regulation of fatty acid oxidation. Beside the correlation with BMI, Adiponectin level is associated with the Insulin-Resistance and so also linked with Type II Diabetes. Adiponectin is associated also with glucose- und lipometabolism.


The formerly proposed diagnostic value of the high molecular weight form of adiponectin was not verified using a commercially available test system for the determination of HMW adiponectin. Blueher et al. clearly demonstrate that regarding the diagnosis of insulin resistance, measured by whole body glucose uptake below  $40 \mu\text{mol}/\text{kg} \cdot \text{min}$ , total adiponectin as determined with this test kit, is with an area of 0.92 under the receiver-operating curve, of greater diagnostic value.

Furthermore it is involved in inflammatory processes and therewith it is of importance for appearance of arteriosclerosis and coronaritis, thus the determination of Adiponectin level in plasma could serve to estimate the risk of coronary disease. Beside this Adiponectin influences further physiological processes as for example the angiogenesis.

Research Area

Cancer kit; Cell Biology and Cellular Response kit; Developmental Biology kit; Metabolism kit; Neuroscience kit; Signaling Transduction kit

## Images



ARG80823 Human Adiponectin ELISA Kit example of standard curve image

ARG80823 Human Adiponectin ELISA Kit results of a typical standard run with optical density reading at 450 nm.