

ARG81540 Human FCN2 ELISA Kit

Package: 96 wells
Store at: 4°C

Component

Cat. No.	Component Name	Package	Temp
ARG81540-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81540-002	Standard	2 X 50 ng/vial	4°C
ARG81540-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81540-004	Antibody conjugate concentrate (100X)	1 vial (100 µl)	4°C
ARG81540-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81540-006	HRP-Streptavidin concentrate (100X)	1 vial (100 µl)	4°C
ARG81540-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81540-008	25X Wash buffer	20 ml	4°C
ARG81540-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81540-010	STOP solution	10 ml (Ready to use)	4°C
ARG81540-011	Plate sealer	4 strips	Room temperature

Summary

Product Description	ARG81540 Human FCN2 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human FCN2 in serum, plasma (heparin) and cell culture supernatants.
Tested Reactivity	Hu
Tested Application	ELISA
Specificity	There is no detectable cross-reactivity with other relevant proteins.
Target Name	FCN2
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	0.39 ng/ml
Sample Type	Serum, plasma (heparin) and cell culture supernatants.
Standard Range	0.78 - 50 ng/ml
Sample Volume	100 µl

Precision	Intra-Assay CV: 6.0% Inter-Assay CV: 7.0%
Alternate Names	Ficolin-B; Hucolin; Ficolin-beta; Collagen/fibrinogen domain-containing protein 2; 37 kDa elastin-binding protein; Ficolin-2; EBP-37; ficolin-2; Serum lectin p35; P35; L-ficolin; FCNL

Application Instructions

Assay Time	~ 5 hours
------------	-----------


Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	FCN2
Gene Full Name	ficolin (collagen/fibrinogen domain containing lectin) 2
Background	The product of this gene belongs to the ficolin family of proteins. This family is characterized by the presence of a leader peptide, a short N-terminal segment, followed by a collagen-like region, and a C-terminal fibrinogen-like domain. This gene is predominantly expressed in the liver, and has been shown to have carbohydrate binding and opsonic activities. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]
Function	May function in innate immunity through activation of the lectin complement pathway. Calcium-dependent and GlcNAc-binding lectin. Enhances phagocytosis of <i>S.typhimurium</i> by neutrophils, suggesting an opsonic effect via the collagen region. [UniProt]
Highlight	Related products: FCN2 antibodies ; FCN2 ELISA Kits ; New ELISA data calculation tool: Simplify the ELISA analysis by GainData

Images



ARG81540 Human FCN2 ELISA Kit standard curve image

ARG81540 Human FCN2 ELISA Kit results of a typical standard run with optical density reading at 450 nm.