

ARG81589 Human RSPO4 ELISA Kit

Package: 96 wells
Store at: 4°C

Component

Cat. No.	Component Name	Package	Temp
ARG81589-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81589-002	Standard	2 X 10 ng/vial	4°C
ARG81589-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81589-004	Antibody conjugate concentrate (100X)	1 vial (100 µl)	4°C
ARG81589-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81589-006	HRP-Streptavidin concentrate (100X)	1 vial (100 µl)	4°C
ARG81589-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81589-008	25X Wash buffer	20 ml	4°C
ARG81589-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81589-010	STOP solution	10 ml (Ready to use)	4°C
ARG81589-011	Plate sealer	4 strips	Room temperature

Summary

Product Description	ARG81589 Human RSPO4 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human RSPO4 in serum, plasma (heparin, EDTA) and cell culture supernatants.
Tested Reactivity	Hu
Tested Application	ELISA
Specificity	There is no detectable cross-reactivity with other relevant proteins.
Target Name	RSPO4
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	15.6 pg/ml
Sample Type	Serum, plasma (heparin, EDTA) and cell culture supernatants.
Standard Range	31.2 - 2000 pg/ml
Sample Volume	100 µl

Precision	Intra-Assay CV: 5.8% Inter-Assay CV: 6.6%
Alternate Names	CRISTIN4; Roof plate-specific spondin-4; hRspo4; R-spondin-4; C20orf182

Application Instructions


Assay Time	~ 5 hours
------------	-----------

Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	RSPO4
Gene Full Name	R-spondin 4
Background	This gene encodes a member of the R-spondin family of proteins that share a common domain organization consisting of a signal peptide, cysteine-rich/furin-like domain, thrombospondin domain and a C-terminal basic region. The encoded protein may be involved in activation of Wnt/beta-catenin signaling pathways. Mutations in this gene are associated with anonychia congenital. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Sep 2009]
Function	Activator of the canonical Wnt signaling pathway by acting as a ligand for LGR4-6 receptors. Upon binding to LGR4-6 (LGR4, LGR5 or LGR6), LGR4-6 associate with phosphorylated LRP6 and frizzled receptors that are activated by extracellular Wnt receptors, triggering the canonical Wnt signaling pathway to increase expression of target genes. Also regulates the canonical Wnt/beta-catenin-dependent pathway and non-canonical Wnt signaling by acting as an inhibitor of ZNRF3, an important regulator of the Wnt signaling pathway. [UniProt]
Highlight	Related products: New ELISA data calculation tool: Simplify the ELISA analysis by GainData
PTM	Tyr-112 may be phosphorylated; however as this position is probably extracellular, the vivo relevance is not proven. [UniProt]



ARG81589 Human RSPO4 ELISA Kit standard curve image

ARG81589 Human RSPO4 ELISA Kit results of a typical standard run with optical density reading at 450 nm.