

# Product datasheet

info@arigobio.com

# ARG82652 Mouse PDGF BB ELISA Kit

Package: 96 wells Store at: 4°C

# Component

| Cat. No.     | Component Name                        | Package              | Temp  |
|--------------|---------------------------------------|----------------------|---|
| ARG82652-001 | Antibody-coated microplate            | 8 X 12 strips        | 4°C. Unused strips should be sealed tightly in the air-tight pouch. |
| ARG82652-002 | Standard                              | 2 X 10 ng/vial       | 4°C   |
| ARG82652-003 | Standard/Sample<br>diluent            | 30 ml (Ready to use) | 4°C   |
| ARG82652-004 | Antibody conjugate concentrate (100X) | 1 vial (100 μl)      | 4°C   |
| ARG82652-005 | Antibody diluent<br>buffer            | 12 ml (Ready to use) | 4°C   |
| ARG82652-006 | HRP-Streptavidin concentrate (100X)   | 1 vial (100 μl)      | 4°C   |
| ARG82652-007 | HRP-Streptavidin diluent buffer       | 12 ml (Ready to use) | 4°C   |
| ARG82652-008 | 25X Wash buffer                       | 20 ml                | 4°C   |
| ARG82652-009 | TMB substrate                         | 10 ml (Ready to use) | 4°C (Protect from light)  |
| ARG82652-010 | STOP solution                         | 10 ml (Ready to use) | 4°C   |
| ARG82652-011 | Plate sealer                          | 4 strips             | Room temperature  |
|              |                                       |                      |   |

## Summary

| Product Description | ARG82652 Mouse PDGF BB ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse PDGF BB in serum, plasma (EDTA, heparin) and cell culture supernatants. |  |  |  |
|---------------------|---|--|--|--|
| Tested Reactivity   | Ms  |  |  |  |
| Tested Application  | ELISA   |  |  |  |
| Target Name         | PDGF BB   |  |  |  |
| Conjugation         | HRP   |  |  |  |
| Conjugation Note    | Substrate: TMB and read at 450 nm.  |  |  |  |
| Sensitivity         | 32 pg/ml  |  |  |  |
| Sample Type         | Serum, plasma (EDTA, heparin) and cell culture supernatants.  |  |  |  |
| Standard Range      | 62.5 - 4000 pg/ml   |  |  |  |
| Sample Volume       | 100 μΙ  |  |  |  |
| Precision           | Intra-Assay CV: 5.4%<br>Inter-Assay CV: 5.8%  |  |  |  |

Alternate Names

SIS; Becaplermin; Platelet-derived growth factor subunit B; IBGC5; PDGF subunit B; c-sis; SSV; PDGF-2; Proto-oncogene c-Sis; Platelet-derived growth factor beta polypeptide; PDGF2; Platelet-derived growth factor B chain

## **Application Instructions**

**Assay Time** 

~ 5 hours

#### **Properties**

Form

96 well

Storage instruction

Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note

For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol

**PDGFB** 

Gene Full Name

platelet-derived growth factor beta polypeptide

Background

This gene encodes a member of the protein family comprised of both platelet-derived growth factors (PDGF) and vascular endothelial growth factors (VEGF). The encoded preproprotein is proteolytically processed to generate platelet-derived growth factor subunit B, which can homodimerize, or alternatively, heterodimerize with the related platelet-derived growth factor subunit A. These proteins bind and activate PDGF receptor tyrosine kinases, which play a role in a wide range of developmental processes. Mutations in this gene are associated with meningioma. Reciprocal translocations between chromosomes 22 and 17, at sites where this gene and that for collagen type 1, alpha 1 are located, are associated with dermatofibrosarcoma protuberans, a rare skin tumor. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2015]

Function

Growth factor that plays an essential role in the regulation of embryonic development, cell proliferation, cell migration, survival and chemotaxis. Potent mitogen for cells of mesenchymal origin (PubMed:26599395). Required for normal proliferation and recruitment of pericytes and vascular smooth muscle cells in the central nervous system, skin, lung, heart and placenta. Required for normal blood vessel development, and for normal development of kidney glomeruli. Plays an important role in wound healing. Signaling is modulated by the formation of heterodimers with PDGFA (By similarity). [UniProt]

Highlight

Related products:


PDGF BB antibodies; PDGF BB ELISA Kits;

Related news:

The role of HDGF in tumor angiogenesis New ELISA data calculation tool: Simplify the ELISA analysis by GainData

Cellular Localization

Secreted. Note=Released by platelets upon wounding. [UniProt]



## ARG82652 Mouse PDGF BB ELISA Kit standard curve image

ARG82652 Mouse PDGF BB ELISA Kit results of a typical standard run with optical density reading at 450 nm.