

ARG83796 Human RANK Ligand / TNFSF11 / TRANCE ELISA Kit

Package: 96 wells
Store at: 4°C, -20°C

Summary

Product Description	ARG83796 Human RANK Ligand / TNFSF11 / TRANCE ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human RANK Ligand / TNFSF11 / TRANCE in serum, plasma and cell culture supernatants.
Tested Reactivity	Hu
Tested Application	ELISA
Target Name	RANK Ligand / TNFSF11 / TRANCE
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	31.25 pg/ml
Sample Type	Serum, Plasma and Cell culture supernatants.
Standard Range	62.5 – 4000 pg/ml
Sample Volume	50 µl
Alternate Names	TRANCE; Osteoprotegerin ligand; CD254; sOdf; Receptor activator of nuclear factor kappa-B ligand; OPTB2; RANKL; OPGL; Tumor necrosis factor ligand superfamily member 11; hRANKL2; TNF-related activation-induced cytokine; Osteoclast differentiation factor; ODF; CD antigen CD254

Application Instructions

Assay Time	4 hours
------------	---------

Properties

Form	96 well
Storage instruction	Store the kit at 4°C, -20°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	TNFSF11
Gene Full Name	tumor necrosis factor (ligand) superfamily, member 11
Background	This gene encodes a member of the tumor necrosis factor (TNF) cytokine family which is a ligand for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. This protein was shown to be a dendritic cell survival factor and is involved in the regulation of T cell-dependent immune response. T cell activation was reported to induce expression of this gene and lead to an increase of osteoclastogenesis and bone loss. This protein was shown to activate antiapoptotic kinase AKT/PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor-associated factor (TRAF) 6, which indicated this protein may have a role in the regulation of cell

apoptosis. Targeted disruption of the related gene in mice led to severe osteopetrosis and a lack of osteoclasts. The deficient mice exhibited defects in early differentiation of T and B lymphocytes, and failed to form lobulo-alveolar mammary structures during pregnancy. Two alternatively spliced transcript variants have been found. [provided by RefSeq, Jul 2008]

Function Cytokine that binds to TNFRSF11B/OPG and to TNFRSF11A/RANK. Osteoclast differentiation and activation factor. Augments the ability of dendritic cells to stimulate naive T-cell proliferation. May be an important regulator of interactions between T-cells and dendritic cells and may play a role in the regulation of the T-cell-dependent immune response. May also play an important role in enhanced bone-resorption in humoral hypercalcemia of malignancy. [UniProt]

Images



ARG83796 Human RANK Ligand / TNFSF11 / TRANCE ELISA Kit standard curve image

ARG83796 Human RANK Ligand / TNFSF11 / TRANCE ELISA Kit results of a typical standard run with optical density reading at 450 nm.